

OVIPOSITION AND FEEDING ALLELOCHEMICALS FOR

CHILO PARTELLUS

FROM SORGHUM AND MAIZE CULTIVARS;

BIOASSAY, ISOLATION AND CHARACTERIZATION:

L. C. I. P. E. LIBRARY
ACC. No 97-9660
CLASS No ^{THE} 595.729 KIB
AUTHOR KIBWANA, RICHARD
TITLE OVIPOSITION AND

BY

BOGITA RICHARD KIBWAGE.


A thesis submitted in **partial fulfilment** for the degree of

Master of Science in the University of Nairobi.

1994.

DECLARATION:

This Thesis is my original work and has not been presented
for a degree in any other University.




BOGITA RICHARD KIBWAGE.

This Thesis has been submitted for examination
with our approval as Supervisors.



PROF. P. M. GITU.
CHEMISTRY DEPARTMENT,
UNIVERSITY OF NAIROBI.



PROF. K. N. SAXENA.
THE INTERNATIONAL CENTRE
FOR INSECT PHYSIOLOGY AND
ECOLOGY (I.C.I.P.E).

DEDICATION:

To my father, *E. Bogita*; mother, *B. Nyarinda*; brother, *J. Nyagaka*
and sisters, *Mary, Josephine, Janice, Zipporah* and *Ester*.

ACKNOWLEDGEMENTS:

I wish to thank most sincerely **Prof. K. N. Saxena and Prof. P.M. Gitu, my supervisors**, for the deep interest they took in this work, for the immense advice, encouragement and guidance they gave me during the course of this work and for critically reading the entire manuscript. I wish to thank **Dr. W. Lwande, Prof. A. Hassanali and Mr. Nyarango** for their timely and continual support and advice.

I wish to thank the director of the **I.C.I.P.E** Prof. T. R. Odhiambo for allowing me to use facilities at the I.C.I.P.E for the study. The part played by the staff at the **B.C.E.R.U, C.P.R.U** and the Insectary units of the I.C.I.P.E is gratefully acknowledged.

I am grateful to **N.A.B.S.A** who paid for the MS and NMR spectra and to Mr. Wanyama (I.C.I.P.E) and Prof. E. Dagne (Addis Ababa University) for running the MS and NMR spectra respectively.

My deep appreciation is expressed to my **parents, Mr and Mrs. Bogita, my Uncle and aunt, Mr. and Mrs. Mokaya**, for their material and financial support in the course of this work.

Finally I wish to thank Mr. J. Maosa for teaching me the computer packages I used to type out and print this thesis and Miss I. Ombaki for allowing me to use her P.C to write out this thesis.

| TABLE OF CONTENTS: | PAGE: |
|--|--------------|
| LIST OF TABLES, GRAPHS AND FIGURES: | vii |
| ABSTRACT: | ix |
| CHAPTER 1 : | 1 |
| INTRODUCTION: | 1 |
| 1.0.0 General Considerations: | 1 |
| 1.1.0 The Pest <i>Chilo Partellus</i> | 3 |
| 1.1.1 Description and Biology: | 3 |
| 1.1.2 Plant Infestation: | 4 |
| 1.1.3 Control of the Pest: | 5 |
| 1.2.0 The Host - Pest Relationship: | 6 |
| 1.2.1 Factors that Influence Oviposition and Feeding of <i>Chilo Partellus</i> | 7 |
| 1.2.2 Host Plant Resistance: | 9 |
| 1.2.3 Mechanisms of Host Plant Resistance: | 10 |
| 1.3.0 Allelochemicals: | 11 |
| 1.3.1 The Chemistry of Allelochemicals: | 12 |
| 1.3.2 Bioactivity of Allelochemicals: | 16 |
| 1.3.3 Insect Feeding Deterrents: | 18 |
| 1.4.0 Previous Work on Maize, Sorghum, Insect Pests, Oviposition and Feeding: | 19 |
| 1.4.1 Previous Work on Maize and Sorghum: | 20 |
| 1.4.2 Previous Work on Insect Pest Oviposition and Feeding: | 23 |
| 1.5.0 Objectives Of The Study: | 26 |

| | |
|---|----|
| CHAPTER 2: | 27 |
| RESULTS AND DISCUSSION: | 27 |
| 2.1 Yields of Extracts From the Sorghum and Maize Whorls: | 28 |
| 2.2 Sorghum; Oviposition Bioassay: | 30 |
| 2.3 Compounds Identified From IS 1044 Pet-ether Extract: | 39 |
| 2.4 Differences Between the Sorghum Cultivars: | 54 |
| 2.5 Maize; Feeding Bioassay: | 56 |
| 2.6 Compounds Identified From the MP 704 Ethyl Acetate Extract: | 58 |
| 2.7 Differences Between the Maize Cultivars: | 66 |
| 2.8 Comparison Between the Sorghum and Maize Cultivar Allelochemicals: | 66 |
| 2.9 DISCUSSION. | 67 |
| CHAPTER 3 | 73 |
| EXPERIMENTAL: | 73 |
| 3.0 General Experimental Procedures: | 73 |
| 3.1 Chromatography: | 74 |
| 3.2 Biological Activity Tests (Bioassays): | 75 |
| 3.3 Plant Material: | 76 |
| 3.4 Insects: | 76 |
| 3.5 Extraction: | 77 |
| 3.6.0 Fractionation and Isolation: | 77 |
| 3.6.1 Fractionation and Isolation of IS 1044 Pet-ether Extract Compounds: | 78 |
| 3.6.2 Isolation and Purification of MP 704 Ethyl Acetate Extract Compounds: | 78 |
| 4.0 REFERENCES: | 79 |

| LIST OF TABLES, GRAPHS AND FIGURES: | | PAGE. |
|--|------|---|
| Tables | 1.0 | Classes of Allelochemicals. 11. |
| | 1.1 | Examples of Reported Allelochemicals. 12. |
| | 1.2 | Some Reported Allelochemical Compounds from Sorghum and Maize. 20. |
| | 1.3 | Some Sorghum Feeding / Oviposition Allelochemicals. 24. |
| | 2.1 | Extracts Yields from Sorghum and Maize Whorls. 28. |
| | 2.2 | Average Yield per Dry Weight Material. 29. |
| | 2.3 | Percentage Oviposition Preference. 30. |
| | 2.4 | Percentage Insects Laying $B \leq 1.5T$ 32. |
| | 2.5 | Average Eggs on Treated / Blank Surfaces. 33. |
| | 2.6 | <i>n</i> -Alkanes from the Petroleum ether Extract. 40. |
| | 2.7 | The <i>n</i> -alkanes Mass Spectra Summary Table. 41. |
| | 2.8 | The Cultivar Relative Feeding Response (Maize). 56. |
| Graph | 2.1a | Cultivar Oviposition Response (Sorghum). 36. |
| | 2.1b | % Insects with O.P. ≤ -20 37. |
| | 2.4a | Cultivar Feeding Response (maize). 57. |
| Fig. | 2.2a | Cultivar Petroleum Ether GC. Profiles (Sorghum). 43. |
| | 2.2b | <i>n</i> -Alkanes Mass Spectra. 45. |
| | 2.2c | Mass Spectrum of the Isolated Methyl-ester. 50. |
| | 2.2d | Proton n.m.r Spectrum of the Methyl-ester. 51. |
| | 2.2e | ¹³ C-n.m.r Spectrum of the Methyl-ester. 52. |

| | | |
|------|---|-----|
| 2.2f | Infrared Spectrum of the Methyl-ester. | 53. |
| 2.3a | HPLC Profiles of the Sorghum Cultivars. | 55. |
| 2.5a | Cultivar HPLC Profiles (Maize). | 62. |
| 2.5b | Mass Spectra of the Isolated Compounds I, II & IV. | 63. |

ABSTRACT:

A study was carried out to isolate and identify the chemicals from sorghum and maize that determine the oviposition and feeding responses, respectively, of the moth and larvae of *Chilo partellus* a serious pest of these crops. Studies were carried out on a resistant and a susceptible cultivar of maize (MP 704 resistant; INBRED A susceptible) as well as sorghum (IS 1044 resistant; IS 18363 susceptible). The allelochemicals were extracted sequentially with petroleum ether, ethyl acetate and methanol. The sorghum cultivars showed quantitative rather than qualitative differences while the maize cultivars showed both qualitative and quantitative differences in respect to the concentrations of the constituent chemical compounds.

The maize feeding bioassay was carried out using impregnated cellulose acetate discs. The sorghum ovipositional bioassay was carried out using filter paper as plant substrate. The maize methanolic extracts showed greater feeding stimulation than ethyl acetate extracts and these in turn showed greater feeding stimulation than the petroleum ether extracts. The susceptible cultivar extracts elicited more feeding stimulation than those of the resistant cultivars. The sorghum petroleum ether extracts showed significant *C. partellus* ovipositional inhibition unlike the ethyl acetate and methanol extracts which showed very weak ovipositional activity or none at all. The pet-ether extract of the resistant cultivar showed the greatest oviposition inhibition.

Chemical study of the extracts revealed a variety of compounds on the leaf whorls. A series of *n*-alkanes with carbon chain C_{10} - C_{18} were common in all the maize and sorghum cultivars. These were however found to have no observable effect on *Chilo partellus* oviposition. The compound, **methyl 11,14,17-eicosatrienoate**, isolated from the sorghum

resistant cultivar petroleum ether extract showed significant ovipositional inhibition. This compound was in large amounts in the resistant cultivar. Three compounds were isolated from the maize resistant cultivar ethyl acetate extract; **4-hydroxybenzoic acid**, **4-ethoxybenzoic acid** and **4,5,6,7-tetrahydro-2-1H-isoindole-1,3(2H)-dione**. 4-Hydroxybenzoic acid was in large amounts in the susceptible cultivar compared with the resistant cultivar and is a known *C. partellus* feeding stimulant. 4-Ethoxybenzoic acid, also in larger amounts in the susceptible cultivar compared to the resistant, has not been tested independently for bioactivity. 4,5,6,7-Tetrahydro-2-1H-isoindole-1,3(2H)-dione was found in large amounts in the resistant cultivar. It has not been tested independently for bioactivity although a study of the H.P.L.C profiles of the cultivars suggests that this compound, along with another that was difficult to isolate, could be responsible for the observed reduced *C. partellus* feeding stimulation in the resistant cultivar by acting as antifeedants.

Structural elucidation of each of these compounds was performed on the basis of their spectroscopic data. The range of compounds isolated and characterized indicates that there may be a vast reservoir of yet untapped allelochemicals in food-crops which could be useful in the breeding of pest resistant cultivars of the principle food-crops maize and sorghum.

CHAPTER 1 :

INTRODUCTION:

1.0 General considerations:

In the tropics, maize and sorghum are some of the principle food crops. They are also used as fodder, fuel and building material. Generally grain yields in farmers fields in Africa and India are low. One of the several factors held responsible for the low yields is losses to the insect pests (Seshu, 1982). The stem borer, *Chilo partellus*, is one of the most widespread and serious maize and sorghum pests in the Indian continent, East, Central and Southern Africa (Young and Teete, 1977; Van Hamburg, 1980; Alighali, 1985). The damage caused to the grains by this pest ranges between 24-36% in maize and 15-45% in sorghum (Seshu Reddy and Sum, 1991; Seshu, 1988 and Alighali, 1986).


To date, a series of maize and sorghum cultivars produced by national and international breeding programs have been evaluated for resistance and varieties have been identified with different levels of preference for oviposition, feeding and establishment of *Chilo partellus* larvae (Jotwani, 1981; Seshu, R.K.V. 1985; Singh and Raina *et al.*, 1989). The degree of susceptibility of the cultivar depends on the successful larval establishment, amount of feeding and development (Alighali and Saxena, 1988).

Host plant resistance is a complex phenomenon involving multiple factors and plays a major role in the control of pest population in crops. One of the factors that plays a vital

role in the location and colonization of crops by pests is the presence or absence of secondary plant metabolites. The production of secondary plant metabolites has been shown to be mainly responsible for the resistance in different crops, for example in rice (Sogawa and Pathak, 1970), sorghum (Woodhead,1983; Haskin and Gorz,1985; Torto *et al.*,1990) and in wheat (Niemeyer, 1989). Factors which may be of major importance in relation to one pest or disease may be secondary or not relevant in another.

It has been suggested that the resistance to *Chilo partellus* by the host plant may be due to a combination of chemical compounds thus, green leaf alcohols, aldehydes, terpenes, etc. that are found on the leaf surface wax (Ampofo and Nyangiri, 1986; Bernays et al, 1983; Woodhead and Janeja, 1987). Indeed, 2,4-dihydroxy-8-methoxy-1,4-benzoxazin-3-one (DIMBOA), (I), found in young maize leaves, is regarded as one of the secondary plant compounds responsible for maize resistance to *Ostrinia nubilalis*, (Robinson *et al.*, 1978) though it does not appear to be a factor in the response of *Chilo partellus* to maize.

It is now well documented that the different behavioral responses shown by an insect, which lead to the selection of and successful establishment on its host plant, are influenced in part, and often predominantly, by allelochemicals. It is, therefore, important and essential to investigate the allelochemical bases for the different colonization of a pest insect on different host cultivars, to provide a better understanding of the underlying principles. Work relating to the *C. partellus* feeding allelochemicals was initiated at the International Centre of Insect Physiology and Ecology (I.C.I.P.E) in 1985 (Torto *et al.* 1990). This research sought to study *C. partellus* oviposition and feeding allelochemicals in sorghum and maize, respectively.



(I)

1.1.0 The pest *Chilo Partellus*, (Swinhoe)

(Pyralidae: Lepidoptera)

1.1.1 Description and Biology:

The *Chilo partellus* moth is medium sized and straw coloured. A female lays nearly 500 eggs in masses of 10-80 on the undersurface of a leaf often near the midrib. The eggs are flattish and oval and tend to overlap like fish scales. The eggs hatch in 4-5 days.

Chilo partellus moths have the greater part of their activity during the night. They have nocturnal habits. Emergence of male and female moths peaks 2 to 3 and 6 to 9 hours after onset of night, respectively. Maximum mating takes place during the first night after emergence, declining markedly during the successive nights. Mating commences after midnight reaching a peak between 5 am and 7 am and then declines. Oviposition is maximum during the first night after mating and declines on successive nights. The most suitable period

for oviposition is between 4 pm and midnight (Ramachandran and Saxena, 1991; Kumar and Saxena, 1985).

The larval period lasts from 19 to 27 days, pupation takes place inside the stem and the larvae emerge in 7 to 10 days through the larvae's entry holes. During the dry season the instar larvae enter into diapause which is broken during the rainy season and pupation takes place giving rise to the first generation of adults.

Forty one species of the genus *Chilo* are known. Twenty five of these infest cereals and eighteen species occur in Africa (Beever, *et al.*, 1990). Essentially it is a pest in hot lowland area and is seldom found above an altitude of 1500 meters (Hill , 1983).

1.1.2 Plant infestation:

The main hosts of *Chilo partellus* larvae are maize, sorghum, sugar-cane, rice, millet and wheat. The larvae may attack various parts of the crop at various stages of the plant growth; however, the larvae prefer feeding on the young plant rather than older ones (Sigh and Raina, 1989; Teete *et al.*, 1983).


In the field, the female moth of *Chilo partellus* lays eggs mostly on the leave blades of maize and sorghum. The newly emerged larvae, after wondering around, finally enter into the whorls of young maize and sorghum plants. Once in the whorls, they feed on leaves for a while and cause a characteristic pattern of small holes and scarification of leaf epidermis. Later on, they feed on the growing stem of the young plants resulting in dead hearts. After third moult, they bore into the stem and start tunnelling. The fully mature larvae pupate inside the plant tissue. Depending on the prevailing temperature and relative humidity the incubation, larval and pupal period vary considerably from 2-9, 15- 41 and 2-12 days, respectively. A


complete generation takes 4-7 weeks. In Kenya oviposition by *Chilo partellus* starts 2-3 weeks after germination and peaks 2-3 weeks latter (Alighali, 1985).

1.1.3 Control of the pest:

Several parasites and predators are known to suppress pest density; however, there use in *Chilo partellus* control has not been fully successful. Ploughing up and destroying the stubble after harvesting is strongly recommended. Early planting with a high seed rate and removal of affected plants is advantageous (Seshu Reddy, 1981). Chemical control with Endosulfan (II) or Carboryl (III) is fairly effective but is not environmental friendly.

Chilo species are particularly difficult to control largely because of the cryptic and nocturnal habits of the adult moths and the protection offered by the stem or cob of the host crop to the developing stages (Sigh and Raina, 1989). Insecticidal measures have proved difficult to apply effectively. In the case of sugar cane, high crop density makes access for ground spaying almost impossible, while in rice cultivation insecticide application is detrimental to beneficial insects and can cause serious pollution of water ways and fish (Beevor *et al.*, 1990). Other alternative control measures that would overcome the drawbacks of insecticides are therefore required.





(III)

The use of sex pheromones is being tried as one of the environmental friendly measures of chilo control (Beevor, *et al.*, 1990; *Ibid.* 1977; and Unnithan and Saxena 1991). Feeding and oviposition deterrents would be desirable when environmental factors are considered; moreover, feeding and oviposition deterrents and stimulants from plants or crops provide knowledge about the interrelationship between insect pests and plants. This in turn gives possibilities for developing other methods of crop protection by breeding crops of insect resistant varieties.

1.2.0 The Host - Pest relationship:

The survival and development of a pest, and consequently, the damage on the plant (crop) is directly related to the interaction between the insect pest and the host plant; thus the expression of cultivar resistance and/or insect preference for feeding, oviposition and orientation are typical of the pest-host relationship.

Various plant cultivars differ in their levels of susceptibility or resistance to the insect pest (Jotwani *et al.*, 1987; Jotwani and Davis, 1980; Lal and Pant, 1980; Jotwani, 1981; Singh *et al.*, 1983; Dabrowski and Kidiavai, 1983). The factors that determine the level of resistance

/ susceptibility have been arranged into two broad categories by Saxena (1969 *ibid.* 1985): (1) the insect colonizing responses, leading to the establishment of its population on the plant and (2) the plant characters which determine these response. The colonizing responses were distinguished by Saxena (1985) into the following main categories; (1) orientation of the insect determining its arrival/arrest on, or avoidance of a plant, (2) feeding, (3) utilization of ingested food determining the insect's nutrition, (4) Development of the larvae, (5) egg production (fecundity) in the adult and (6) oviposition.

The lower the insects response in each of these categories to a cultivar, the greater will be the plant's resistance and thus, the less the insect's preference for that cultivar. Ideally these lead to reducing the insect's three major behavioral responses:- oviposition orientation and/or feeding.

1.2.1 Factors that influence oviposition and feeding of *chilo partellus*.

There are possibly many factors that influence the oviposition and feeding of *Chilo partellus* and other plant pests on the maize and sorghum cultivars, however, these can be categorized into two factors, namely;

(a) Bio-physical properties of the host plant:

Ampofo (1985) reported that the choice of oviposition site is probably influenced by the leaf surface trichomes, thus the lower leaf surface and the mid-rib concavity (the smooth areas of the plant) were noted to be preferred for oviposition with up to 92% preference for 3-4 weeks old maize plants. Ampofo suggested that tactile stimuli due to increased trichome density on the leaf surface, probably combined with other factors may be an important factor on *Chilo partellus* selection of oviposition site and orientation on the host plant. Kumar and

Saxena (1985) reported that the *Chilo partellus* moth prefers glass, polythene sheet, wax paper then filter paper in this order for the ovipositing site, while Raina (1982) reported that sorghum sootfly (*Atherigona soccata*) prefer green colour to white and a surface with ridges to a smooth substrate for oviposition.

(b) Bio-chemical properties of the host plant:

Kira *et al.* (1968) noted considerable increase in *Ostrina nubilalis* (hubner) egg production when moths were provided drinking water while Derridj and Fiala (1983) observed a direct relationship between the concentration of soluble sugars in the foliage of different maize hybrids and the number of *O. nubilalis* egg masses laid on them. Differences in glucose and/or fructose concentrations were implicated in the selection of site by *O. nubilalis*. Ampofo (1985) observed that plant exudates from ICZ2-CM maize lines significantly increased oviposition above that of distilled water while exudates from INBRED A and especially ICZ1-CM depressed oviposition. The exudates were observed to be of different chemical composition which therefore affected *Chilo partellus* oviposition and longevity differently. It has been shown that feeding by third instar larvae of *Chilo partellus* on sorghum bicolor is mediated by several surface and tissue chemicals (Torto *et al.*, 1990, *ibid.* 1991).

Considering the feeding response , for example, insect feeding involves various behavioral responses which can be divided into four distinct steps (Dethier, 1966; Munakata, 1977; Schoonhoven 1982; Miller and Sticker, 1984); (1) host plant recognition and orientation, comprising locomotion which brings the insect to it's food and cessation of locomotion on arrival, also termed as arrest; (2) initiation of feeding (biting, probing or sucking); (3) continuation of feeding and (4) termination of feeding as a result of satiation.

It has been suggested that the acceptance or rejection of a plant as food by an insect is determined by a number of factors which include the intensity of olfactory and gustatory

feeding stimulants, the intensity of repellents and deterrents, the metabolic state of the insect and learning acquired as a result of previous feeding experience (Dethier, 1982).

Electrophysiological studies at the sensory level show that chemoreceptors are the means by which the insect detects plant chemicals and is able to differentiate between a stimulant and a deterrent compound thus insects are said to respond to a summation of inputs (gestalt) from various chemoreceptor which perceive complex mixtures of compounds present in their food (Schoonhoven and Jermy, 1977; Dethier, 1980; Stadler, 1982).

1.2.2 Host plant resistance:

The seriousness of the damage caused by *Chilo partellus* larvae and other insect pests on host plants and their effect on grain or crop yield has prompted research concentrated on screening experiments with a view to identifying and developing resistant varieties. Efforts to identify sources of resistance to *Chilo partellus* and to incorporate the resistance into agronomically acceptable maize and sorghum cultivars are in progress in many research institutions.

Four factors known to affect the expression and stability of resistance of a plant cultivar are genetic, environmental, biochemical and morphological factors (Sigh, 1978, Ampofo, and Nyangiri, 1986).

It is the biochemical factors, however that are of interest to this research. Every green plant is inherently resistant to some phytophagous insect. This is to a certain extent attributed to the chemicals present in the plant. Therefore plants and especially resistant cultivars may develop defence chemistry to enable them to survive in a given habitat.

The chemical resistant factor in a plant (allelochemicals) to a large extent affect the behavioral and metabolic processes of the insect pest. Several chemicals have been identified which are essential or prohibitive to the development of the insect pest (Smith *et al.*,1979; Sagawa and Pathak, ; Torto *et al.*, 1990; Woodhead and Bernays, 1987). The improper balance of these allelochemicals in a cultivar can change the susceptibility level of the cultivar (Auclair *et al.*, 1957; Maxwel *et al.*, 1976; Maltais, 1951; Ampofo and Nyangiri, 1986).

1.2.3 Mechanisms of host plant resistance:

While it is clear that certain plant cultivars possess specific morphological and chemical characteristics which can be advantageously used for insect control, the mechanisms of resistance are generally complex and possibly interrelated (Jotwani, 1981; Painter, 1951).

Observed mechanisms include;

- (i) Antibiosis; affecting insect establishment, survival, development and egg production on host plants.
- (ii) Preference/non-preference for oviposition, feeding and shelter (Painter, 1958) also referred to as antoxenosis by Kogen and Ortman (1978).
- (iii) Tolerance in plants involving repair regeneration of their damaged tissues (Kalode and Pant, 1966; Jatwani, 1981).

1.3.0 Allelochemicals:

Plant allelochemicals are secondary plant products that have a significant effect to the organism or plant of a species different from its source. Allelochemicals could be subdivided into four classes according to the effect on the receiving and the emitting species, where the allelochemical is the substance produced or acquired by the plant that when it contacts another species in the natural context thereby evokes in the receiver the response shown (see table 1.0 below).

Table 1.0 Classes of allelochemicals.

| Allelochemical | Evoked physiological reaction |
|----------------|---|
| i) Kairomone | adaptively favourable to the receiver but not to the emitter. |
| ii) Allomone | adaptively favourable to the emitter but not to the receiver. |
| iii) Synomone | adaptively favourable to both the emitter and the receiver. |
| iv) Apneumone | adaptively favourable to the receiving organism but detrimental to an organism of another species that may be found in or on the non living material. |

1.3.1 The chemistry of allelochemicals:




Allelochemicals belong to a few major classes of chemical compounds. These include terpenoids, phenolic compounds, phenylpropane derivatives, flavonoids, long chain cyanides,


alkaloids, purines and steroids. However, simple chemical compounds such as alkanes, alcohols, aldehydes and fatty acids have also been reported to show activity as allelochemicals. Examples of reported allelochemicals are listed in table 1.1 below.

Table 1.1 Examples of reported allelochemicals.


| Allelochemical and chemical class: | Source and reported activity: | Author/reference |
|--|---|---|
| <p>Phenolic compounds;</p> <p><i>p</i>-Hydroxybenzoic acids</p> <p><i>p</i>-Hydroxybenzaldehyde</p> <p>Phenolic acids</p> | <p><i>S. bicolor</i>; feeding stimulant for <i>C. partellus</i>.</p> <p><i>S. bicolor</i>; <i>C. partellus</i> feeding stimulant, <i>I. nubilalis</i> feeding inhibitor.</p> <p><i>S. bicolor</i> as non-deterrent esters but released as acids deterring <i>I. migratoria</i> feeding.</p> | <p><i>Torto et al.</i> 1990.</p> <p><i>Torto et al.</i>, 1990;</p> <p>Woodhead, 1982.</p> <p>Woodhead and Bernays 1978.</p> |
| <p>Alkaloids and nitrogen compounds;</p> <p>N-acyl analog of nicotine (IV)</p> <p>Non-protein analogue of phenylalanine and tyrosine <i>eg.</i> Hiptagin (V).</p> | <p><i>N. rapenda</i> plant; toxic to tobacco horn worm (<i>manducasexta</i>).</p> <p>Laguminous pasture plant (<i>I. pendunculatus</i>) roots and foliage; feeding deterrent of grass-grab (<i>Cosalytra zealandica</i>).</p> | <p>Huesing and Jones 1987.</p> <p>Granderam and Sutherland 1986.</p> |

| Allelochemical and chemical class: | Source and reported activity: | Author /reference |
|--|--|--|
| Tannins | American deciduous tree defence against insect feeding. | Barbosa and Kirschner 1987. |
| Terpenoids Momordicine II (VI) | Cucurbit (<i>M. charantia</i>) reduces feeding of red pumpkin beetle (<i>Aulacophora foveicollis</i>) | Chandravadana 1987. |
| Tenulin (VII) | Bitter weed (<i>H. amarum</i>) Antifeedant and reduces growth and development of European corn borer <i>O. nubilalis</i> | Broughtton <i>et al.</i> 1986 |
| Flavonoids Luteolin-7-O-(6-O- melanylglucoside) (VIII) | Carrot leaf; oviposition stimulant to swallowtail moth (<i>P. polyxenes</i>). | Feeny 1988. |
| Hydrocarbons 32-carbon chain hydrocarbon. | S. bicolor; cultivar resistance to stem borers. | Woodhead and Janeja 1987. |
| Short chain hydro-carbons C ₁₀ to C ₁₈ . | Rice cultivar IR-22 Resistance to Brown plant hopper (<i>N. lugens</i>) | Woodhead and Padgham 1988. |
| Alcohols, aldehydes and fatty acids. | 1). S. bicolor; feeding deterrents for <i>L. migratoria</i> 2). Rice; resistance to <i>N. lugens</i> . | Woodhead 1983. Woodhead and Padgham 1988. |





(VII)



(VIII)

1.3.2 Bioactivity of allelochemicals:

Plant allelochemicals can determine the establishment of an insect species on the plant by determining its orientation, feeding and/or oviposition behaviour towards the plant and/or influencing the metabolism serving as;

- a) Nutrients promoting normal metabolic processes resulting in the insects survival, development and egg production.
- b) toxins interfering with the metabolic processes causing failure of the insect's survival, development and egg production (Saxena, 1985 *ibid.* 1987).

The allelochemical may provide olfactory stimuli perceivable at a distance from the plant or contact stimuli perceivable after the insect's arrival on the plant. The orientation of the insect can be determined by the olfactory stimuli whereas its feeding and oviposition responses can be determined by the contact and to some extent by the olfactory stimuli (Dethier, 1947; Thorseinson, 1966; Norris and Kogen, 1980; Pathak and Dole, 1983).


The action of allelochemicals may be influenced by a single dominant allelochemical or different sets of allelochemicals acting together or in opposition (Dethier, 1982; Miller and Sticker, 1984; Woodhead and Bernice, 1978).

While many plant allelochemicals may be directly synthesized and are found in the plant in the bioactive form, in the environment for which they are produced, some allelochemicals have been reported that are produced *de novo* (damage induced).

In the resistance of sorghum bicolor, for example, to attack by *L. migratoria*, the relative unpalatability of seedling is the result of several deterrent chemicals which are produced only at the time of feeding. This is true of HCN, stored as the glucoside Dhurrin, and the phenolic acids, stored as esters. As a result of damage to the plant tissues these

substances are brought into contact with enzymes to produce the bioactive compounds though the substrate themselves are not bioactive (Woodhead and Bernice, 1978)

The enzymatic hydrolysis of Dhurrin to *p*-hydroxybenzaldehyde and HCN is stoichiometric as outlined below.



In maize, damage induced DIMBOA production protects maize against stalk borers (Cooper Driver and Wain, 1976; Bernice *et al.*, 1977).


1.3.3 Insect feeding deterrents:

Plant chemicals which possess feeding deterrent activities against insects are known as antifeedants (Dethier, *et al.* 1966). They prevent or decrease the feeding of the insect without killing it. The insect remains near or on the treated plant and dies from starvation (Dethier, 1980; Munakata, 1977). It has been suggested that an antifeedant is concerned with deterring the initiation of feeding therefore may act as a resistance factor protecting plants against insect attack (Munakata, 1977). However, it is plausible that in some cases it is the insect that has evolved the means of detecting and avoiding a potentially deleterious compound. Many deterrents are also toxic to insects.

Plant deterrent chemicals are derived from virtually all classes of natural products. However studies to date suggest that most potent insect antifeedants belong to the terpenoid and alkaloid classes (Schoonhoven, 1982).

Among the terpenes, the tetranortriterpenoid azadirachtin (IX) isolated from the neem tree *Azadirachta Indica Meliaceae* (Butterworth *et al.* 1972) is an antifeedant for a host of different lepidopteran, coleopteran and orthopteran insects.

While many antifeedants are insect species specific, broad spectrum antifeedants are also known; apart from azadirachtin, the mustard glycoside sinigrin, the carbohydrate phlorizin, the flavonoid glycoside rutin, several quinones, naphthoquinones and some ellagitaninins have been found to be broad spectrum insect antifeedants (Schoonhoven, 1972; Bernays and Chapman, 1977; Hedin *et al.*, 1977; Norris, 1986; Jones and Klocke 1987).



(IX)

1.4.0 Previous work on maize, sorghum, insect pests, oviposition and feeding:

Extensive research has been done on sorghum and maize regarding their biology, cultivation and breeding for pest resistance on host plant-pest relationship as reflected by the references quoted; however, not much has been done on the biochemical aspects of the plants in relation to their insect pests, especially on the chemicals possibly involved the oviposition and feeding of *Chilo partellus*. Outlined below is some of the previous work.

1.4.1 Previous work on maize and sorghum:

Tabulated below (table 1.2) are some of the compounds and their reported activities that have been isolated from maize and sorghum but may not be necessarily involved in *C. partellus* oviposition and/or feeding bioactivity.


Table 1.2. Some reported allelochemical compounds from sorghum and maize.

| Compound: | Source and reported bioactivity | Author/reference |
|---|---|--------------------------------|
| C_{27} , C_{29} and C_{31} n-alkanes, dotriacontanal, <i>n</i> -octacosanal, <i>n</i> -tricosanal, <i>n</i> -dotriacotanol, <i>n</i> -octacosol, <i>n</i> -tricosanol, several esterified primary alcohols and fatty acids. | Epicuticular wax of two maize seedlings gl2 and GL (no bioassay done) | Bianchi <i>et al.</i> 1975. |
| (z)-Hexenyl acetate, (z)-hexenol, cyclosalirene (sesquiterpene), x-ylangene, | Corn leaf volatile (no bioassay done) | Buttery and Lousa 1984. |
| 2-heptanone, 4-hepten-2-one, and caryophyllene. | Corn leaf volatile (no bioassay done) | Buttery and Lousa 1984. |


| Compound: | Source and reported bioactivity | Author/reference |
|--|---|-----------------------------------|
| Toluene, hexanal, (z)-3-hexen-1-ol, <i>m</i> -xylene, <i>o</i> -xylene, (z)-3- hexen-1-ol acetate, nonanal and decanal. | <i>S. bicolor</i> seedlings (no bioassay done) | Lwande and Bentley 1987. |
| (+)Taxifolin-7- <i>o</i> - β - glucoside,(Xa), eriodictyol-5- <i>o</i> - β -glucoside (Xb), 4,2',4',6',-tetra- hydroxychalcone-4- <i>o</i> - β -glucoside (Xc), 4-procyanidins (XIa - XIId), | Sorghum grains | Rene <i>et al</i> 1986. |
| Flavan-4-ols | Sorghum grains, leaf, and tissue (of resistant cultivar to grain mold) | Ramamurthi <i>et al.</i> 1986. |
| β -Caryophyllene, longifolene, bazzanene, cyclosativene and <i>x</i> -ylangene (sesquiterpenes). | Volatile of cone roots (no bioassay done) | Buttery and Loisa, 1985. |

Most of the above compounds vary in concentration with the cultivar of the plant and with the plant maturation (Buttery and Louisa, 1984; Woodhead, 1982). While most of the above compounds have not been tested for bioactivity, some of them could be playing a role in the insect pest behaviour.


str.Xn - Xld




(Xa)



(Xb)




(Xc)



(XIa), R = H.

(XIb), R = Glucose.



(XIc), R' = H.

(XId), R' = Glucose.

1.4.2 Previous work on insect pest oviposition and feeding:

Besides the compounds reported elsewhere in this thesis, the following are some of the compounds that have been reported to have allelochemical effects on insect pest oviposition and/or feeding:

Table 1.3 Some sorghum feeding/oviposition allelochemicals.

| Compound | source and reported bioactivity | Author and reference |
|--|--|---|
| Dhurrin, <i>p</i> -Hydroxybenzaldehyde, <i>p</i> -hydroxybenzoic acid, and procyanidin (unidentified) | <i>S. bicolor</i> ; feeding deterrents to green aphids | Dreyer <i>et al.</i> 1980. |
| Sugars; sucrose, glucose, fructose and xylose, <i>p</i> -hydroxybenzaldehyde, <i>p</i> -hydroxybenzoic acid, <i>p</i> -hydroxycinnamic acid, <i>p</i> -hydroxy- <i>m</i> -methoxy cinnamic acid and several phenolic analogs. | <i>S. bicolor</i> ; feeding stimulants to <i>C. partellus</i> larvae. | Torto <i>et al.</i> 1990 <i>ibid.</i> 1991. |

Raina (1981) reported oviposition deterrence of sorghum shootfly (*Atherigonia soccata*) by what he associated to an unidentified deterrent pheromone in the water soluble glue with which females attach their eggs to the leaves. Under low population density conditions, the females lay one egg per sorghum plant, hence repeated oviposition is deterred by the suggested pheromone.

Unnithan *et al.* (1987) and Saxena (1990) reported that acetone extracts of resistant, susceptible and wild sorghum elicited high ovipositional response to shootfly on maize, a non host plant, therefore suggesting a potential method for reducing the population of the pest by diverting oviposition to non host plants. However, the oviposition stimulants in the extract have not yet been identified.

Kumar (1986) reported that the maize plants infested with *Chilo partellus* larvae elicited more oviposition than the uninfested plants. Ampofo (1985) and Ampofo and Nyangiri (1986) reported differing oviposition among maize genotypes with INBRED A eliciting maximum oviposition. They suggested that certain conditioning stimuli were central to larvae acceptance or rejection of the plant on which eggs were laid. Their analysis indicated that different chemical characteristics played a certain role in larvae acceptance of the plant. Kumar (1985) made similar observations on the ovipositional responses to the susceptible and resistant sorghum cultivars.

Saxena (1987) employing techniques specially developed to test the role of cultivars in oviposition reported remarkable differences in oviposition of the susceptible, IS 18363, and the resistant, IS 1044, cultivars. The number of eggs laid as well as the ovipositional preference were high for IS 18363, while the number of eggs as well as the ovipositional preference were low and medium respectively for IS 1044. Other workers (I.C.I.P.E annual report 1988) show that *n*-hexane and/or acetone extracts of susceptible and resistant cultivars of sorghum (IS 18363, IS 1044) exhibit deferent levels of oviposition stimulation and/or inhibition.

1.5 Objectives of the study:

The review of the feeding and ovipositional allelochemical work on the spotted borer *C. partellus* shows that no detailed and methodical study has been carried out to identify the chemicals from sorghum and maize which affect the oviposition and feeding, respectively, of the moth and larvae of this insect; Besides, the literature available on this subject suggests that not much work has been undertaken to demonstrate whether or not there are any allelochemical bases for oviposition and feeding preferences shown by the moths and larvae to different sorghum and maize cultivars .

The objectives of the study therefore are;

- I). Isolation of allelochemicals of maize and sorghum through sequential solvent extraction of the leaves followed by fractionation of the extracts using chromatographic techniques and their structural elucidation using spectroscopic methods.
- II). Carrying out feeding and oviposition bioactivity assays of the crude extracts and where possible of the isolated compounds against the stalk borer *C. partellus* moths and larvae.

Ultimately this study would enhance the knowledge on *C. partellus* oviposition, feeding and establishment on the host plants sorghum and maize. This would therefore lead to better methods of control of the pest; methods that would be environmental-friendly and both technologically and socio-economically feasible for the small scale resource poor farmer.

CHAPTER 2:

2.0. RESULTS AND DISCUSSION:

The results of this work are outlined below. The differences between the resistant cultivars and the susceptible cultivars of both maize and sorghum are clearly depicted. Consequently an attempt is made to use the observed cultivar chemical differences to explain the observed biological behaviour of the *Chilo partellus* pest on the host crops. As reflected by the quotation below, this results serve to demonstrate the important linkage between chemistry and biology in the natural world.

"Much of life can be understood in rational terms if expressed in the language of chemistry. It is an international language, a language for all time, a language that explains where We have come from, what We are and where the physical world will allow us to go. Chemical language has great aesthetic beauty and links the physical sciences to the biological sciences. Unfortunately, the full use of this language to understand life processes is hindered by a gulf that separates chemistry from biology. The gulf is not nearly so wide as the one between humanities and sciences. Yet chemistry and biology are two distinct cultures and the rift between them is serious, generally unappreciated and counterproductive....." (*Nobel Laureate Arthur Kornberg on Chemistry and Biology*).

The yields of the extracts from the cultivar whorls and the oviposition and feeding bioactivity of the crude extracts and some of the isolated and characterized compounds are reported. The characterized compounds include a series of *n*-alkanes of carbon chain length C₁₀ to C₁₈, three phenolic compounds, one methyl-ester and one heterocyclic compound.

2.1 Yields of extracts from the sorghum and maize whorls:

Table 2.1 Yields of the extracts.

| Cultivar | Plant material | | Weight of extract | | | Total Yield |
|---|-----------------|----------------|-------------------|---------------|---------------|-------------|
| | Fresh wt gms | Dry wt. gms | Pet Ether mgs. | EtAc. mgs. | MeOH. mgs. | |
| Sorghum season I (28th June 1992): | | | | | | |
| IS1044 | 443.01 | 49.1 | 422.03 | 451.63 | 2217.64 | 3091.3 |
| IS18520 | 551.03 | 56.1 | 233.1 | 613.79 | 3023.52 | 3870.41 |
| IS18363 | 482.83 | 9.3 | 279.24 | 385.18 | 2694.64 | 3359.06 |
| Sorghum season II (8th September 1993): | | | | | | |
| IS1044 | 750.23 | 63.2 | 795.42 | 756.42 | 3714.27 | 5266.11 |
| IS18520 | 459.68 | 46.8 | 489.83 | 869.10 | 4281.83 | 5640.76 |
| IS18363 | 357.47 | 36.5 | 471.05 | 693.50 | 4853.61 | 6018.16 |
| Maize season I (June 1991): | | | | | | |
| MP 704 | 674 | 71.22 | 214.6 | 136.0 | 4895.2 | 5245.8 |
| INBRED A | 800 | 80.36 | 382.0 | 219.9 | 15000.0 | 15601.9 |
| Maize season II (28th June 1992): | | | | | | |
| MP 704 | 335.97 | 35.5 | 214.64 | 354.39 | 4286.25 | 4855.28 |
| INBRED A | 722.72 | 72.6 | 318.71 | 563.31 | 8058.77 | 8940.79 |
| Maize season III (8th September 1993): | | | | | | |
| MP 704 | 579.19 | 61.2 | 765.80 | 1124.6 | 10206.55 | 12096.95 |
| INBRED A | 706.79 | 71.0 | 707.94 | 922.01 | 9894.41 | 11524.36 |

Table 2.2 Average yield in mgs per gm of dry weight plant material (mg/g).

| Cultivar | Pet-ether | Ethyl acetate | methanol | total |
|----------|-----------|---------------|----------|--------|
| Sorghum. | | | | |
| IS 1044 | 10.6 | 10.59 | 51.97 | 73.14 |
| IS 18520 | 7.34 | 14.76 | 72.70 | 94.76 |
| IS 18363 | 6.82 | 13.41 | 93.82 | 116.51 |
| Maize. | | | | |
| MP 704 | 7.19 | 10.09 | 118.75 | 136.03 |
| INBRED A | 6.37 | 7.83 | 145.67 | 159.87 |

The yields of the petroleum ether (P.E.), ethyl acetate (EtAc.) and methanol (MeOH) extracts were dependent on the cultivar, the season of planting and on the age at harvesting time (tables 2.1 & 2.2). In general the susceptible cultivars of both sorghum and maize gave higher yields of the total extracts compared to the resistant cultivars. This was generally true of both the ethyl acetate and methanol extracts for sorghum and methanol extract for maize, however for sorghum there was a remarkably higher yield of the petroleum ether extract from the resistant (IS 1044) cultivar compared to the susceptible (IS 18363), while for maize there was higher yield of the ethyl acetate extract from the resistant cultivar (MP 704) compared to the susceptible (INBRED A).

2.2 Sorghum; oviposition bioassay:

The relative suitability of the test extracts and samples was compared on the basis of the ovipositional preference (O.P) for the cultivar (extract) relative to the non-plant substrate (blank). The O.P. represented the percentage as $100(T-B)/(T+B)$, T and B being the number of eggs laid on the treated half disc and the blank half disc, respectively. Positive O.P values reflect attraction and/or contact stimulation while negative values reflect, avoidance and/or contact inhibition of the oviposition by the cultivar extract or sample. O.P values close to zero reflect lack of involvement of the plant extract or sample in oviposition.

Table 2.3 Percentage oviposition preference. (O.P \pm Std Error)

| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|---|----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----|
| IS 1044 P.E. | 13.35 ± 11.88 | -22.32 ± 7.59 | -35.4 ± 10.22 | -31.31 ± 7.77 | -31.78 ± 11.14 | |
| IS 18363 P.E. | 16.31 ± 13.23 | -11.59 ± 15.17 | -5.82 ± 15.76 | -27.1 ± 10.39 | -29.04 ± 14.78 | |
| IS 18520 P.E. | 18.04 ± 12.94 | -18.81 ± 12.46 | -24.68 ± 17.85 | -30.02 ± 12.92 | -29.00 ± 12.12 | |
| IS 1044 EtAc. | | 20.98 ± 11.82 | -8.51 ± 14.49 | 1.42 ± 16.55 | -0.95 ± 19.5 | |

| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|---|----------------------|-----------------------|-----------------------|-----------------------|----------------------|----------------------|
| IS 18363 EtAc. | | 16.96 ± 13.86 | 21.81 ± 10.88 | -15.56 ± 13.87 | | |
| IS 1044 MeOH | | -7.74 ± 11.57 | 7.78 ± 14.72 | 9.43 ± 8.58 | 14.4 ± 14.18 | |
| IS 1044 P.E.:- | | | | | | |
| Remixed Fxn | | -11.96 ± 10.21 | -16.85 ± 10.78 | -21.76 ± 12.02 | | |
| Fxn. I | | | -7.61 ± 18.61 | -2.72 ± 8.48 | 9.66 ± 14.57 | 29.71 ± 15.18 |
| Fxn II. | | | 18.0 ± 9.74 | 22.07 ± 12.55 | | |
| Fxn III. | | -20.56 ± 13.31 | -24.56 ± 11.13 | -27.40 ± 11.31 | | |
| <i>p</i> -Hydroxy - benzaldehyde | 0.39 ± 12.43 | -1.34 ± 9.57 | 16.97 ± 11.58 | 8.06 ± 12.25 | -0.14 ± 14.00 | |
| <i>p</i> -Hydroxybenzoic acid | -3.83 ± 12.00 | -2.00 ± 11.98 | 16.97 ± 12.99 | -4.32 ± 9.48 | 6.03 ± 10.06 | |

Table 2.4 Percentage insects laying more than 1.5 times as many eggs on the blank as on the treated surface.

| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|---|-------|-------|-------|-------|-------|-----|
| IS 1044 P.E. | 50 | 45 | 60 | 75 | 75 | |
| IS 18363 P.E. | 16.7 | 42.86 | 35.7 | 66.6 | 68.75 | |
| IS 18520 P.E. | | 33.3 | 50 | 58 | 58 | |
| IS 1044 EtAc. | 33.3 | 16.7 | 41.67 | 33.33 | 41.67 | |
| IS 18363 EtAc. | | 14 | 21.43 | 41.67 | | |
| IS 1044 MeOH | | 33 | 30.77 | 23.1 | 25 | |
| IS 1044 P.E.:- | | | | | | |
| Remixed Fxn | | 53.3 | 42.86 | 66.67 | | |
| Fxn. I | | | 41.67 | 37.5 | 27.8 | 33 |
| Fxn II. | | | 25 | 25 | | |
| Fxn III. | | 50 | 55.56 | 60.14 | | |
| <i>p</i> -Hydroxy- benzaldehyde | 33.3 | 33.3 | 22.2 | 27.27 | 333 | |
| <i>p</i> -Hydroxybenzoic acid | 30.77 | 35.71 | 27.78 | 47.37 | 27.78 | |

Table 2.5 Average number of eggs laid on the treated compared to the blank surfaces.


| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|---|-------------|------------|------------|------------|------------|-----|
| IS 1044 P.E. | 101 | 105 | 76 | 91 | 83 | |
| | ± 16.7 | ± 15.5 | ± 15.9 | ± 13.6 | ± 12.4 | |
| Blank. | 134 | 138 | 147 | 160 | 174 | |
| | ± 12.21 | ± 30.3 | ± 18.1 | ± 17.1 | ± 25.9 | |
| IS 18363 P.E. | 120 | 127 | 85 | 106 | 77 | |
| | ± 21.0 | ± 22.2 | ± 17.2 | ± 17.1 | ± 16.5 | |
| Blank. | 113 | 184 | 122 | 186 | 137 | |
| | ± 25.4 | ± 27.4 | ± 27.4 | ± 20.3 | ± 19.9 | |
| IS 18520 P.E. | | 97 | 77 | 87 | 77 | |
| | | ± 20.4 | ± 19.1 | ± 17.8 | ± 11.0 | |
| Blank. | | 133 | 144 | 158 | 154 | |
| | | ± 17.9 | ± 28.8 | ± 20.0 | ± 21.4 | |
| IS 1044 EtAc. | 146 | 209 | 139 | 144 | 95 | |
| | ± 20.0 | ± 27.8 | ± 26.9 | ± 25.4 | ± 24.8 | |
| Blank. | 127 | 126 | 174 | 156 | 134 | |
| | ± 25.6 | ± 19.2 | ± 23.5 | ± 29.6 | ± 27.6 | |
| IS 18363 EtAc. | | 163 | 161 | 114 | | |
| | | ± 23.3 | ± 24.9 | ± 24.8 | | |
| Blank. | | 102 | 98 | 123 | | |
| | | ± 15.6 | ± 16.9 | ± 17.3 | | |

| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|---|----|------------|------------|------------|------------|------------|
| IS 1044 MeOH | | 120 | 141 | 195 | 174 | |
| | | ± 19.7 | ± 19.7 | ± 22.6 | ± 24.0 | |
| Blank. | | 147 | 120 | 141 | 111 | |
| | | ± 21.1 | ± 18.3 | ± 16.4 | ± 17.8 | |
| IS 1044 P.E:- | | | | | | |
| Remixed Fxn | | 139 | 122 | 124 | | |
| | | ± 20.0 | ± 19.1 | ± 20.4 | | |
| Blank. | | 186 | 158 | 181 | | |
| | | ± 24.4 | ± 20.3 | ± 23.0 | | |
| Fxn. I | | | 105 | 133 | 115 | 132 |
| | | | ± 25.1 | ± 14.0 | ± 17.9 | ± 19.2 |
| Blank. | | | 93 | 141 | 92 | 119 |
| | | | ± 23.8 | ± 13.7 | ± 18.8 | ± 27.1 |
| Fxn II. | | | 194 | 162 | | |
| | | | ± 26.6 | ± 20.4 | | |
| Blank. | | | 149 | 110 | | |
| | | | ± 32.4 | ± 22.6 | | |
| Fxn III. | | 79 | 78 | 84 | | |
| | | ± 15.0 | ± 11.7 | ± 14.9 | | |
| Blank. | | 108 | 107 | 150 | | |
| | | ± 18.8 | ± 11.7 | ± 18.5 | | |


| Sample/Dose ($\mu\text{g}/\text{cm}^2$) | 10 | 20 | 40 | 80 | 160 | 320 |
|--|-------------------|-------------------|-------------------|-------------------|-------------------|-----|
| <i>p</i> - H y d r o x y - benzaldehyde | 122 ± 16.0 | 130 ± 17.4 | 134 ± 18.0 | 159 ± 22.7 | 151 ± 23.5 | |
| Blank. | 148 ± 23.1 | 125 ± 15.0 | 103 ± 18.9 | 139 ± 20.0 | 134 ± 22.4 | |
| <i>p</i> -Hydroxybenzoic acid | 148 ± 23.5 | 152 ± 21.5 | 144 ± 22.1 | 116 ± 14.9 | 119 ± 14.7 | |
| Blank. | 150 ± 19.4 | 149 ± 19.0 | 114 ± 23.8 | 130 ± 17.7 | 121 ± 16.7 | |

These oviposition bioassay results are summarized in the graphs 2.1a and 2.1b below.

Graph 2.1a Cultivar Oviposition Response (sorghum).



% Insects Laying B \geq 1.5T.



Graph 2.1b % Insects with O.P. \leq -20.

a) bioassay of the crude extracts:

As shown on the tables 2.3 and 2.4, graphs (2.1a) and (2.1b) the petroleum ether extracts showed the greatest oviposition inhibition of *Chilo partellus* moth. The resistant cultivar (IS 1044) extract showed the greatest inhibition. At high doses (80 μ l/cm² and higher) the activity of all the pet-ether extracts is very high and remains relatively unaffected by the increase in concentration. At lower doses, however, the activity increased with the concentration. The ethyl acetate extracts show relatively weak oviposition stimulation of *C. partellus* moth at low concentrations and the susceptible (IS 18363) showed slightly higher stimulation than the resistant (IS 1044). The methanol extracts seemed not to have any effect on the oviposition of *C. partellus* moth at all the doses tested.

From table 2.5 it is noted that for the pet-ether extracts, the number of eggs laid on the blank is significantly higher than the number laid on the treated as compared to the ethyl acetate and methanol extracts. Bioassay done with petroleum ether on the treated surface and hexane on the blank show that the pet-ether on its own has no effect on the *C. partellus* oviposition. The insects showed no ovipositional preference for pet-ether nor for hexane. The average no of eggs laid on the treated and the blank showed no significant difference (T; 143 & B; 132).

b) Isolated fractions and authentic samples:

The qualitative oviposition bioassay of the three fractions from the column showed that fraction I and II, individually, have relatively very low oviposition stimulation activity. Thus, on their own, they are not responsible for the observed inhibition in the parent crude extract. Fraction III, however, showed relatively high oviposition inhibition though not as high as the parent crude extract (Tables 2.3, 2.4 and 2.5). Comparing the activity of the crude extract, the

column fraction III and the remixed fractions, it was noted that the bioactivity of the remixed fractions was lower than that of the crude extract and that of fraction III, however, higher than that of the ethyl acetate and methanol crude extracts.

p-Hydroxybenzaldehyde and *p*-hydroxybenzoic acid which have been isolated from the sorghum ethyl acetate extract (Torto *et al.* 1990) were assayed and they showed relatively weak oviposition stimulation at low doses but decreased in activity at higher doses.

2.3 Compounds identified from IS 1044 pet-ether extract:

Fraction I from the column was analyzed by GC-MS and identification was confirmed by co-injection in the GC with the authentic samples. A series of *n*-alkanes of C₁₀ to C₁₈ were characterized as outlined in table 2.6 below.

As would be expected, the GC profile (fig. 2.2a) shows the characteristic arrangement of the peaks at regular intervals which decrease slightly from one peak to the next one. The observed interval represents the CH₂ difference between successive alkanes.

The concentrations of the *n*-alkanes were notably varying with the season of planting/harvesting. During season (I), pentadecane, hexadecane and heptadecane were most abundant while in season (II) dodecane, tridecane and tetradecane were most abundant (fig.2.2a).

Table 2.6 *n*-alkanes characterized from the pet-ether extract.

| <i>n</i> -alkanes | GC min. | Rt. | % comp. | % comp. | MASS- SPECT. | m/e of M+ ion | % M+ abd. |
|-------------------|------------|-----|---------|---------|-----------------|------------------|--------------|
| | | | Season | Season | | | |
| | | | I | II | | | |
| Decane | 9.17 | | 0.233 | 2.046 | 142 | 10 | |
| Undecane | 11.272 | | 0.961 | 1.355 | 156 | 10 | |
| Dodecane | 13.085 | | 0.496 | 3.517 | 170 | 8 | |
| Tridecane | 14.69 | | 1.604 | 2.570 | 184 | 9 | |
| Tetradecane | 16.156 | | 4.621 | 4.391 | 198 | 8 | |
| Pentadecane | 17.532 | | 14.836 | 1.028 | 212 | 8 | |
| Hexadecane | 18.823 | | 34.46 | 0.239 | 226 | 8 | |
| Heptadecane | 19.876 | | 18.516 | 0.190 | 240 | 7 | |
| Octadecane | 20.769 | | 1.928 | 0.099 | 254 | 6 | |

As observed from the mass spectra (fig. 2.2b table 2.6 & table 2.7) the molecular ion peak (M⁺) of the *n*-alkanes is always present but decreases in intensity with the *n*-alkane chain length. The fragmentation pattern is characterized by clusters of peaks, and the corresponding peaks of each cluster are 14 (CH₂) mass units apart. The largest peak in each cluster represents a C_nH_{2n+1} fragment. This is accompanied by C_nH_{2n} and C_nH_{2n-1} fragments. The most abundant fragments are at C₃ and C₄ and the fragment abundance decreases in a smooth curve down to M-C₂H₅. The M-CH₃ peak is characteristically very weak or missing.

Table 2.7 Summary of the mass spectra fragmentation pattern of the *n*-alkanes and the relative abundance of the fragments.

| <i>n</i> -alkane chain | | C ₁₀ | C ₁₁ | C ₁₂ | C ₁₃ | C ₁₄ | C ₁₅ | C ₁₆ | C ₁₇ | C ₁₈ |
|--|-----|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|
| Fragment (+). | m/e | | | | | | | | | |
| CH ₃ (CH ₂) ₂ | 43 | 98 | 96 | 80 | 80 | 80 | 78 | 84 | 76 | 74 |
| CH ₃ (CH ₂) ₃ | 57 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |
| CH ₃ (CH ₂) ₄ | 71 | 21 | 23 | 58 | 61 | 62 | 70 | 83 | 75 | 70 |
| CH ₃ (CH ₂) ₅ | 85 | 15 | 15 | 38 | 40 | 44 | 44 | 58 | 50 | 51 |
| CH ₃ (CH ₂) ₆ | 99 | 5 | 4 | 8 | 11 | 12 | 16 | 20 | 19 | 19 |
| CH ₃ (CH ₂) ₇ | 113 | 4 | 2 | 4 | 4 | 5 | 6 | 12 | 11 | 10 |
| CH ₃ (CH ₂) ₈ | 127 | | 1 | 2 | 2 | 3 | 3 | 4 | 4 | 6 |
| CH ₃ (CH ₂) ₉ | 141 | | | 1 | 1 | 3 | 3 | 3 | 3 | 4 |
| CH ₃ (CH ₂) ₁₀ | 155 | | | | 1 | 2 | 2 | 3 | 3 | 3 |
| CH ₃ (CH ₂) ₁₁ | 169 | | | | | 1 | 2 | 2 | 2 | 2 |
| CH ₃ (CH ₂) ₁₂ | 183 | | | | | | 1 | 2 | 2 | 2 |
| CH ₃ (CH ₂) ₁₃ | 197 | | | | | | | 1 | 1 | 2 |
| CH ₃ (CH ₂) ₁₄ | 211 | | | | | | | | 1 | 1 |
| CH ₃ (CH ₂) ₁₅ | 225 | | | | | | | | | 1 |

As observed and expected, n-alkanes containing more than eight carbon atoms show fairly similar spectra; hence identification depends on the molecular ion peak (Silverstein *et al.* 1981).

The presence of branched alkanes is ruled out from the observed spectra because although the spectra of branched chain hydrocarbons are grossly similar to those of the straight chain compounds, the smooth curve of the decreasing intensities is broken by preferred fragmentation at each branch.




Fig. 2.2a (ii). GC. Profile; Sorghum Resistant
(IS 1044) P.E, Season II.




Fig. 2.2a (i). GC. Profile; Sorghum Resistant
(IS 1044) P.E Season I.




Fig. 2.2a (iv). GC. Profile; Maize Resistant (MP 704) F.E. Season II.




Fig. 2.2a (iii) GC. Profile; Sorghum Susceptible (IS 18365) P.E. Season II.

Fig. 2.2b. Mass Spectra of characterized *n*-alkanes.






Fig. 2.2b. Mass Spectra of characterized *n*-alkanes.









Fig. 2.2b. Mass Spectra of characterized *n*-alkanes.



Fraction (III) from the column was predominantly one compound (over 90% by GC analysis). This was purified using a smaller column affording the compound; **Methyl-11,14,17-Eicosatrienoate**.

structure.




Mass spectra; (Fig.2.2c). The molecular ion peak of a methyl ester of a straight chain aliphatic acid is usually distinct. The most characteristic peak is due to the familiar *McLafferty* rearrangement and cleavage one bond removed from the C=O group. It is clear, however, from the observed spectra that the fragmentation pattern of the compound does not arise from such simple cleavages. There possibly occurs multiple cleavages and rearrangement. These would be favoured by the allylic bonds in the compound. There is observed as would be expected hydrocarbon clusters at intervals of fourteen mass units. Comparing the sample spectra with the computer matched library spectra of the authentic compound it is observed that they are identical.

Proton nuclear magnetic resonance (${}^1\text{H}$ n.m.r) spectrum (Fig. 2.2d) shows resonance at δ 0.90 ppm appearing as a triplet for the three methyl protons of C-20, the three methyl protons of C-1 show resonance at δ 1.30 ppm, the resonance at δ 1.35 ppm is due to the 14 methylene protons of C-3 to C-9 (with overlapping coupling). The resonance of the six olefinic protons of C-11, 12, 14, 15, 17 and 18 occur at δ 5.35 ppm as a triplet, the resonance of the two methylene protons of C-2 at α -position to the C=O group occurs at δ 2.80 ppm as a tetralet. The four equivalent methylene protons of C-13 & C-16 give resonance at δ 2.35 ppm as a triplet while the remaining four methylene protons of C-10 and C-19 show resonance at δ 2.15 ppm as overlapping tetralet and pentatet.

^{13}C n.m.r spectrum (fig. 2.2e) was interpreted by comparing with published ^{13}C n.m.r spectral data (Silverstein, *et al.* 1981) and the assignment was as follows; the ester carbonyl carbon was confirmed by the presence of the resonance at δ 179.01 ppm. The presence of one methyl carbon δ 14.206 ppm, the olefinic carbons (methine) δ 127.212, 127.835, 128.333, 130.265 and 132.009 ppm; thus at least three olefinic bonds. The methylene carbons at δ 33.953, 29.592, 29.094, 27.256, 25.636, 24.731 and 20.588 ppm; at least seven methylene carbons.

Infra-red spectrum (fig. 2.2f) run by preparing the sample as a film cast, confirms the presence of the ester C=O by the strong band at 1710 cm^{-1} corresponding to C=O stretching. The olefinic C-H stretching appears as a weak band at 3224 cm^{-1} . The most characteristic vibrational mode of olefins are the out of plane C-H bending vibration between 1000 cm^{-1} and 650 cm^{-1} these bands are usually the strongest in the spectra of olefins. Therefore the strong band at 725 cm^{-1} confirms the presence of the olefinic bonds. The C=C stretching expected to occur in the region of 1680 cm^{-1} to 1630 cm^{-1} are usually weak and often absent in symmetrical disubstituted or tetrasubstituted trans olefins. This is true of this compound. Olefins which lack symmetry absorb more strongly than trans olefins and internal double bonds generally absorb more weakly than terminal double bonds because of pseudosymmetry. The band at 1260 cm^{-1} corresponds to the asymmetric stretching of the C-O-C bonds and is accompanied with the symmetric stretching at 1075 cm^{-1} . The bands at 2926 cm^{-1} and 2853 cm^{-1} correspond to the asymmetric and symmetric methylene C-H stretching. The weak bands at 1410 cm^{-1} and 1460 cm^{-1} correspond to the methyl C-H symmetric and asymmetric bending. The symmetric band is stronger than the asymmetric band as expected. (Silverstein *et al.* 1981).

Fig. 2.2c. (i) Mass Spectrum of the Isolated Methyl ester.



LIBFITS1#3* x1 Bgd=200 RB281293E +0:00:00
11,14,17-EICOSATRIENOIC ACID, METHYL ESTER
C21.H36.O2.

p844 M966 r844 RFN:55682-88-7
Lib:NBS 31145 Bpk: 79 Mwt: 320

HMR: 65535000
MASS: 79





Fig. 2.2c. (ii) Mass Spectrum of the Authentic sample (Library).

Fig. 2.2d. Proton n.m.r. Spectrum of Isolated Methyl ester.



BSS-23


Fig. 2.2e. ¹³C. n.m.r Spectrum the Isolated Methyl ester.

TOTAL 16
RESOL 78192 -4 HZ
EXREF 77.0000PPM
OES 1896.3745 HZ
NGAIN 13

| NO | FREQ(HZ) | PPM | INT1 |
|----|----------|---------|------|
| 1 | 2574.66 | 132.009 | 1459 |
| 2 | 2935.35 | 130.265 | 2076 |
| 3 | 2891.84 | 128.333 | 2917 |
| 4 | 2860.61 | 127.835 | 1688 |
| 5 | 2866.57 | 127.212 | 1437 |
| 7 | 1767.39 | 78.432 | 5376 |
| 7 | 1735.14 | 77.000 | 9315 |
| 7 | 1703.51 | 75.598 | 5415 |
| 9 | 765.88 | 33.953 | 1902 |
| 10 | 666.81 | 29.592 | 2326 |
| 11 | 655.58 | 29.094 | 5146 |
| 12 | 614.17 | 27.256 | 1836 |
| 13 | 577.67 | 25.636 | 2226 |
| 14 | 557.28 | 24.731 | 1939 |
| 15 | 463.92 | 20.588 | 1388 |
| 16 | 328.84 | 14.286 | 1275 |

Dept

CDM






Fig. 2.2f. Infra red Spectrum of the Isolated Methyl ester.

WASH RECORDING SPECIFY CHART NO. 106508

WAVENUMBER CM⁻¹

RECORDED BY INSTRUMENTS INC., FULLERTON, CALIFORNIA, U.S.A.

PRINTED IN U.S.

2.4 Differences between the sorghum cultivars:


Analysis of the extracts of the three cultivars; IS 1044 (resistant), IS 18520 (tolerant) and IS 18363 (susceptible) by chromatographic techniques, GC and HPLC, showed that the cultivars are different quantitatively rather than qualitatively with respect to the concentrations of the constituent chemical compounds. It is noted from the profiles that all the peaks in the resistant cultivar are present in the tolerant as well as the susceptible cultivar; however,

1) While most of the compounds occur relatively in the same amounts from the GC profile (fig. 2.2a), the compound characterized above with GC retention time 23.115 min occurs with 22.84% in IS 1044 (resistant), 14.03% in IS 18520 (tolerant) but 12.99% in IS 18363 (susceptible). This is apparently the most significant difference in the petroleum ether extract.

2) The most significant difference in the ethyl acetate extracts (HPLC profiles fig.2.3a) is that from the same concentrations of crude extracts the susceptible cultivar (IS 18363) has much higher quantity of compounds corresponding to the peaks at 6.0 min and 15.0 min retention time. These compounds have been identified as *p*-hydroxybenzoic acid and *p*-hydroxybenzaldehyde, respectively (Torto, *et al.* 1990).

3) There is no significant difference in the methanol extracts as all the cultivars show only one peak on analysis by HPLC.

Fig. 2.3a. HPLC Profile of the sorghum cultivars.




2.5 Maize; feeding bioassay:

The *Chilo partellus* larvae feeding response on the crude extracts of the susceptible, INBRED A, and the resistant, MP 704, cultivars were compared on the basis of the calculated relative feeding response X_1/X_0 .

Table 2.8. the relative feeding response of the cultivars crude extracts.

| Cultivar. | Extract conc. $\mu\text{g}/\text{disc}$ | X_1/X_0 Pet- ether | X_1/X_0 Ethyl acetate | X_1/X_0 Methanol |
|-----------|---|----------------------------|-------------------------------|-----------------------|
| Inbred A | 10 | 1.19 | 1.44 | 1.54 |
| | 50 | 1.51 | 1.97 | 2.02 |
| | 100 | 1.79 | 2.69 | 2.43 |
| | 300 | 2.93 | 2.60 | 3.61 |
| | 500 | 2.99 | 3.80 | 4.44 |
| MP 704 | 10 | 1.16 | 1.05 | 1.05 |
| | 50 | 1.36 | 1.30 | 1.42 |
| | 100 | 1.65 | 1.60 | 2.10 |
| | 300 | 2.08 | 2.28 | 2.93 |
| | 500 | 3.36 | 2.90 | 3.41 |

Graph 2.4a Cultivar feeding response (Maize).




In general the methanol extracts were most stimulating followed by the ethyl acetate extracts and lastly petroleum ether extracts. Extracts derived from the susceptible cultivar were more stimulatory than those derived from the resistant cultivars.

In particular, however, for the ethyl acetate extracts, the MP 704 (resistant) response is significantly lower than that of the INBRED A (susceptible) suggesting presence of allomones in the resistant cultivar ethyl acetate extract.

2.6 Compounds identified from the MP 704 ethyl acetate extract:


The compounds of the MP 704 ethyl acetate extract were isolated and purified by Prep. HPLC yielding three compounds corresponding to the peaks I, II, and IV (fig. 2.5a). Compound III could not be obtained in analyzable amounts and it changed colour from white to brown on exposure to air after freeze drying to remove water.

Compound I: 4-hydroxybenzoic acid:




Mass spectra (Fig.2.5b) The molecular ion peak M^+ (138) is large as would be expected of aromatic acids and phenols (Silverstein *et al.* 1981). The other diagnostic peaks that are prominent are $M^+ - OH$ (121) and $M^+ - COOH$ (93). The peak $M^+ - (COOH+CO)$ ie 65 is

predominant and is formed as a result of fragmentation by loss of COOH group followed by rearrangement and loss of CO characteristic of phenols.





The presence and structure of this compound were confirmed by co-injection of the extract with the authentic sample in the H.P.L.C and consequent observation of enhancement.


Compound II: *p*-Ethoxybenzoic acid.




Mass spectra; The parent peak M^+ (166) is large as is expected of aromatic acids. The characteristic peaks are as a result of the fragmentation pattern outlined below (Silverstein *et al.* 1981).



The peak at m/e 121 is obtained from the parent ion by loss of the (COOH) group.




Compound IV: 4,5,6,7-Tetrahydro-2-1H-Isoindole-1,3 (2H)-dione.




Mass spectra; The parent peak, as would be expected, is very intense M^+ (165). The fragmentation pattern may not be a straight forward one, however, cleavage is expected to occur at the C-C bond adjacent to the oxygen atom and at the C-N bonds; Thus leading to

the fragmentation ion peaks outlined below. It is however clear that there occurs rearrangement and multiple fragmentation leading to the other observed peaks.



The structures of the three compounds, I, II and IV, were elucidated by comparison of the sample mass spectrum with the spectra of authentic samples (Silverstein *et al.* 1981). This was done by computerised search and matching of the sample spectrum with the computer library spectra of authentic samples. The computer gave a print out of the five best fitting spectra for further analysis. The five spectra then were manually analyzed peak by peak comparing with the sample spectrum, in conjunction with the expected fragmentation pattern of the various functional groups, to determine the identically fitting spectrum. The isolated sample spectra and the corresponding authentic sample spectra (fig.2.5bi, fig.2.5bii & fig.2.5biii) were found to be identical. The three compounds were therefore easily characterized.

Fig. 2.5a. HPLC profiles of the maize cultivars.



MP 704 (Resistant) - EtAc.

INBRED A (Susceptible) - EtAc.

Fig. 2.5b (i). Mass Spectra of the Isolated compound I.




Fig. 2.5b (i). Mass Spectra of the Authentic sample. (Library).

LIBFITS1#1* x1 Bgd=40 RB281293R +0:00:00
BENZOIC ACID, 4-HYDROXY- p865 M941 r870 RFN:0-00-0
C7.H6.O3. Lib:NBS 5501 Bpk: 121 Mwt: 138

HMR: 65535000
MASS: 121




Fig. 2.5b (ii). Mass Spectra of the Isolated compound II.




Fig. 2.5b (ii). Mass Spectra of the Authentic sample. (Library).

LIBFITS1#1* x1 Bgd=160 RB2812938 +0:00:00

BENZOIC ACID, 4-ETHOXY-
C9.H10.O3.

p581 M745 r582 RFN:619-86-3
Lib:NBS 10762 Bpk: 121 Mwt: 166

HMR: 65535000
MASS: 121




Fig. 2.5b (iii). Mass Spectra of the Isolated compound IV.





Fig. 2.5b (iii). Mass Spectra of the Authentic sample. - (Library).

LIBFITS1#1* x1 Bgd=150 RB2812930 +0:00:00
1H-ISOINDOLE-1,3(2H)-DIONE, 4,5,6,7-TETRAHYDRO-2- p583 M637 r595 RFX:28839-49-8
C9.H11.N.O2. Lib:NBS 10479 Bpk: 165 Mwt: 165

HMR: 65535000
MASS: 165



2.7 Differences between the maize cultivars:

The extracts were analyzed by chromatographic techniques; petroleum ether extracts by GC, ethyl acetate and methanol extracts by HPLC.

1) The petroleum ether extracts of both cultivars showed no obvious qualitative difference; however, a significant difference occurs in the quantity of the compound corresponding to the peak of retention time 23.341 min. The resistant cultivar, MP 704, has 19.27% of the compound while the susceptible, INBRED A, has 7.28. This is the same compound isolated and characterized from sorghum as outlined in section 2.3 above.

2) The ethyl acetate extracts showed remarkable differences qualitatively and quantitatively. As shown by the H.P.L.C profiles (fig. 2.5a) the susceptible cultivar, INBRED A, has a much higher quantity of the compounds of peak I and II; conversely the resistant cultivar, MP 704, has much higher quantities of compounds of peaks III and IV. The other significant difference between the two cultivars is that of the relative amounts of the four compounds in the extract. In the susceptible cultivar the amount of compound I is much higher than that of compound IV, similarly compound II is in higher quantity than IV. In the resistant cultivar, however, the quantity of compound IV is much higher than that of compounds I and II.

3) There was no significant difference in the methanol extracts of the two cultivars as only one peak was observed on analysis by HPLC.

2.8 Comparison between the sorghum and maize cultivar allelochemicals:

A qualitative study of the profiles of the MP 704 and IS 1044 maize and sorghum cultivars reveals some similarities and some differences between the maize and sorghum surface chemistry.

Comparison of the GC profiles (fig. 2.2a) of the pet-ether extracts showed that both maize and sorghum possess the major constituents of the non-polar compounds; the *n*-alkanes from C₁₀ to C₁₈. The compounds with retention times 24.5, 25.224, 26.515, 28.158, 29.49, 30.697, 32.402 and 36.72 minutes, which have not been identified and had no significant effect on the oviposition of *C. partellus*, are common to both maize and sorghum and relatively in similar amounts. The compound of retention time 23.144 minutes, which is most likely the *C. partellus* oviposition inhibitor, occurs in relatively higher quantities in the resistant cultivars of both maize (19%) and sorghum (22%) but in lower quantities in the susceptible cultivars.

The HPLC profiles (fig. 2.2f & fig. 2.5a) show both qualitative and quantitative similarities and differences. The most remarkable similarity is the presence of *p*-hydroxybenzoic acid in relatively large amounts. The sorghum ethyl acetate extract however, is predominantly consisting of *p*-hydroxybenzoic acid and *p*-hydroxybenzaldehyde while the maize extract does not seem to possess detectable amounts of the *p*-hydroxybenzaldehyde, instead it contains compounds II, III, and IV which are interestingly absent in sorghum.

2.9 DISCUSSION.

The available literature on the role of plant characters in determining resistance or susceptibility of different sorghum and maize cultivars to *Chilo partellus* is inadequate. It is the interaction of the different factors rather than any one of them that determine a cultivars resistance or susceptibility to the pest (Saxena, 1990).

These results are in agreement with observations made earlier where larval establishment was found to be high on the leaves of the maize susceptible cultivar INBRED A but low on the resistant cultivar ICZ2-CM. The adult moth oviposition was found to be

high on the leaves of sorghum susceptible cultivar IS 18363 but very low on the resistant cultivar IS 1044 (Ampofo and Nyangiri, 1986; Saxena, 1985 and Saxena, 1987).

The susceptible cultivars of both maize and sorghum give higher quantities of crude extracts than the resistant cultivars, especially so of the methanolic extracts. The methanolic extracts have been shown to consist mainly of sugars which have been shown to be *C. partellus* feeding stimulants acting synergistically with other known stimulants (Torto, *et al.* 1990). This explains the higher feeding stimulation observed in the INBRED A extracts and in the IS 18363 as was observed by Torto, *et al.* It is of importance to note that the sorghum resistant cultivar IS 1044 has higher quantities of the petroleum extractables and more so of the compound methyl 11,14,17-eicosatrienoate compared to the susceptible IS 18363. The maize resistant cultivar MP 704, on the other hand, has higher quantities of the ethyl acetate extractables and especially so of the compounds 4,5,6,7-tetrahydro-2-1H-isoindole-1,3(2H)-dione and another that has not been identified compared to the susceptible cultivar INBRED-A.

It is clear from the bioassay results that the resistant cultivars of the maize and sorghum possess chemicals that inhibit oviposition for sorghum and feeding for maize. From this study it may be concluded that the presence of the compound methyl 11,13,17 eicosatrienoate in high amounts does play a significant role in combination with the alkanes and other unidentified compounds in inhibiting the *C. partellus* oviposition on the highly resistant cultivar IS 1044. It may be concluded too from this study that the presence of the compound 4,5,6,7-tetrahydro-1,3(2H)-dione along with the unidentified compound in high amounts play a significant role in inhibiting (by acting as antifeedants) the feeding of the *C. partellus* larvae on the maize resistant cultivar MP 704. Therefore the low quantities or absence of these compounds in the susceptible cultivar INBRED A leads to high feeding stimulation due to the compound *p*-hydroxybenzoic acid and possibly *p*-ethoxybenzoic acid which are in high quantities in the cultivar. The combination of the various compounds in the

cultivar is clearly important for the cultivars resistance or susceptibility. This is reflected by the feeding bioassay results, the H.P.L.C profiles of the maize cultivars (fig. 2.5a) and by the fact that the various column fractions and the remixed fraction of the sorghum resistant cultivar IS 1044 petroleum extract show much weaker oviposition bioactivity compared to the crude extract.

The bioassay methods used in this study give a reflection of the real life field situation. It however must be noted that the field situation more often than not is far from the laboratory controlled situation, therefore, the improvement of the bioassay conditions so as to approach the field conditions would serve to improve the bioassay results. As observed from the oviposition bioassay there is a wide response variation between individual insects. On any one accession it is not unusual to find that 5% - 10% of the insects do not follow the general pattern. No explanation can be given for this and it is not known whether such insects would consistently be the exceptional ones.

The leaf surface has been shown to be important to locusts and grasshoppers in the selection of food plants (Blaney and Chapman, 1970; Chapman, 1977). Such insects can perceive chemicals in the wax on the leaf surface by contact chemoreceptors on the maxillary palps (Bernays, *et al.*, 1976).

Alkanes are always present in plant surface waxes, but normally the major alkanes have long chains of 27, 29, 31 or 33 carbon atoms (Eglinton and Hamilton, 1963). In sorghum, however, the dominant alkanes vary with cultivar. Bianchi, *et al.*(1978) found C₂₉ and C₃₁ *n*-alkanes predominantly in cultivars SD 102 and Alliance, but Wilkinson and Cummins, (1981) showed that in "bloom" leaves of cultivars Redbine 60 the most abundant *n*-alkanes in the surface wax were C₃₅, C₃₆, C₃₇ and C₃₈, whereas in "bloomless" leaves of the same cultivar, the C₂₅ alkane was in highest concentration.

In young sorghum plants of cultivar CSH 1 and IS 1082, the major alkanes are C₂₇, C₂₉ and C₃₁ with negligible amounts of any alkanes of shorter chain length (Atkin and

Hamilton, 1982). Woodhead, (1983) found in the cultivar 65D surface wax *n*-alkanes of C₁₉ to C₂₅ with C₂₃, C₂₄ and C₂₅ being the most abundant. Woodhead found that only the C₁₉, C₂₁ and C₂₃ *n*-alkanes did reduce feeding of *L. migratoria*.

It is of interest to note that in this study the *n*-alkanes found in the cultivars IS 1044, IS 18520 and IS 18363 are of much shorter carbon chain than any reported earlier, and C₁₄, C₁₅ and C₁₆ are the most abundant in the cultivar IS 1044. Although these *n*-alkanes do not exhibit any observable *C. partellus* oviposition bioactivity for any of the doses tested, they could be acting in combination with the methyl 11,13,17 eicosatrienoate to produce the observed oviposition inhibition in the sorghum resistant cultivar. There is need therefore for a detailed study of the effect of these short chain *n*-alkanes to the pest *C. partellus*.

Alkanes in plant surface waxes have previously been reported to be involved in food selection by *Acyrtosiphon pisum* with C₃₂ serving as a feeding stimulant and by *L. migratoria* with C₁₉, C₂₁ and C₂₃ acting as feeding deterrents (Klingaur *et al.*, 1971; Woodhead, 1983). C₁₀ - C₁₈ *n*-alkanes have been shown to be involved in rice cultivar IR-22 resistance to the brown plant hopper (*N. lugens*) (Woodhead and Padgam, 1988). They have also been reported to affect oviposition of *Delia (Hylemya brassicae)* (Stadler, 1978).

Most higher plant waxes have mainly C₂₀, C₂₂ and C₂₄ acid components of their wax esters (Martin and Juniper, 1970). Seedlings of sorghum cultivar 65D have mainly shorter-chain acid components while seedlings of cultivars CSH 1 and IS 1082 generally have longer-chain fatty acids (Atkin and Hamilton, 1982). Woodhead (1983) reported C₁₂ to C₂₄ esters in the leaf wax of the sorghum cultivar 65D with only traces of odd carbon chain esters.

Methyl esters of C₁₀ and C₁₂ fatty acids have been reported to adversely affect growth and cause mortality in bollworm (Binder *et al.*, 1979), and there are many reports of the use of free short-chain fatty acids as contact insecticides. Mortality was highest with C₁₀ and C₁₂ compounds in, for example, mosquitoes (Saxena and Thorsteinson, 1971) and houseflies (Quraishi, 1971).

The short-chain lauric acid (C₁₂) has been reported to stimulate *L.migratoria* feeding while its methyl ester was strongly deterrent (Woodhead, 1983).

p-Hydroxybenzoic acid, several phenolic compounds and their derivatives have been shown to be *C. partellus* feeding stimulants (Torto *et al.*, 1990 *Ibid.* 1991). Fisk (1990) found that a phenolic extract from sorghum was stimulatory to the homopteran *Peregrinus maidis*, and Baker *et al.* (1986) showed that 4-hydroxybenzaldehyde and some of its derivatives are strong feeding stimulants for the elm bark beetle *Scolytus multistratus*. On the other hand *p*-hydroxybenzaldehyde has been shown to deter feeding by the *L. migratoria* and the aphid *Schizaphis graminum* (Woodhead and Bernays, 1978; Dreyer *et al.*, 1981; Woodhead, 1982).

Not many heterocyclic compounds have been isolated from either sorghum or maize. However, the compound DIMBOA, isolated from young leaves of one resistant cultivar of maize, has been shown to be responsible for maize resistance to *O. nubilalis* and the 1,4-benzoxazin-3-one functional group seemed important for its activity (Robinson *et al.* 1978). It therefore may be of interest to establish the bioactivity of the isolated compound 4,5,6,7-tetrahydro-2-1H-Isoindole-1,3-(2H)-dione and possibly the importance of the isoindole group or the dione arrangement.

The importance of the surface of the young sorghum and maize plant may have been underestimated in previous work when the strongly deterrent nature of the seedlings was attributed to the internal cellular cyanogenic glucoside, dhurrin, and phenolic acid esters (Woodhead and Bernays, 1978). The presence of deterrent chemicals in the surface of the plant is of particular interest in that it provides information to the insect resulting in rejection of the plant prior to or at palpation before the internal feeding deterrents are encountered. As proposed by Chapman (1977) there are obvious advantages to the plant in protecting itself from potential predators by advertising its unsuitability on its surface rather than keeping its defence hidden until it is damaged by being bitten.

From the point of view of a plant breeder, the potential for developing resistant varieties of a crop is probably much higher than may be immediately obvious. The number of possible resistant varieties will be a function of not only of the number of individual chemicals which may individually be increased in amount, but also of the possible combinations and mixtures of individual secondary compounds each of which may be of no obvious significance when tested alone.

CHAPTER 3

(3) EXPERIMENTAL:

3.0 General experimental procedures:

Thin layer chromatography (TLC) was done on MN polygram precoated silica gel/UV254 40x80 mm plates (0.25mm thickness). They were eluted in 10% ethyl acetate - hexane, developed with 50% aqueous sulphuric acid and subsequently heated at 120°C for 30 minutes. Column chromatography was done with (columns 15mm and 10mm diameters) normal phase silica gel (60) particle size 0.040-0.063mm (230-400-mshs ASTM). High performance liquid chromatography (HPLC) was performed on Varian 5000L.C model, equipped with MCH-5 and MCH-10 reverse phase columns. Gas chromatography (GC) profiles were obtained using a Howlett Packard model HP 5890A series gas chromatograph. GC-MS and MS spectra were done on a Howlett Packard model HP 5790A series gas chromatograph coupled to a VG Masslab 12-250 analytical organic mass spectrometer. Nuclear magnetic resonance (proton NMR and ¹³C- NMR) were done on Perkin Elmer Spectrometer model R12B-90MHz. Infra-red spectra was done on a Beckman Infrared spectrophotometer.

The solvents used; hexane, petroleum ether, ethyl acetate, methanol, dichloromethane and acetonitrile were all of analar grade. The hexane used for GC profiles was double distilled while the methanol and acetonitrile used for HPLC were of HPLC grade (solvents were obtained from Aldrich chemical company Ltd, Gillingham UK).

3.1 Chromatography:

Column chromatography was used to isolate compounds from the IS 1044 pet-ether extract. The column was packed using hexane. The extract was dissolved in minimal hexane, introduced in the column and eluted with hexane adjusted in polarity with ethyl acetate and eventually with dichloromethane. The eluates were monitored by their GC profiles.

The ethyl acetate and methanol extracts were analyzed by HPLC. For the profiles, 5mg/ml each of the extracts were prepared in methanol, filtered through a wad of cotton wool plugged in a pasteur pipette and 10 μ l of each extract were injected and analyzed on the MCH-5 (Zorbex 4.6mmx25cm) reverse phase column. The column was eluted with 30% and 20% methanol - water for maize and sorghum extracts, respectively. The flow rate was 1ml per minute, the column pressures were 312 and 270 atm., respectively. The uv. detector, 240nm, was used.

The pet-ether extracts were analyzed by GC. Solution (2 μ l) containing 20 μ g/ μ l of the extract were used. The GC was equipped with a split/splitless injector and a flame ionization detector (FID) at 250°C. The column used was Ultra 1 crosslinked methyl silicone gum of dimension (50mx0.32mm ID; 0.17 μ m film thickness). White spot nitrogen (3.45cm³/sec.) was the carrier gas. Hydrogen (45cm³/sec.) and medical air (360cm³/sec.) were the fuel gases. The total flow was 450cm³/sec. All the GC analyses were done in the splitless mode with the injector temperature at 280°C. The oven temperature was programmed to stay at 45°C for 5min rise to 180°C at a rate of 10°/min and further to 280 at 20°/min The retention times, the peak areas and the percentage composition were calculated on a Howlett Packard model HP 3393A integrator.

3.2 Biological activity tests (bioassays):

Bioassay of the crude extracts, fractions and isolated compounds were done on the *Chilo partellus* larvae and moth to establish the stimulation or inhibition, if any, of feeding and oviposition by the maize and sorghum allelochemicals respectively.

(a) ***Chilo partellus* feeding bioassay;** The no choice feeding tests were conducted in small glass vials (23x27mm) with tight porous caps. A wad of moist cotton wool placed inside the cap maintained a high humidity. Test samples in solvents were applied topically to both sides of the cellulose acetate disks (12mm diameter, 7.902 ± 0.348 mg, Cole- parmer Co. Chicago, Illinois) and were dried in a stream of warm air (Doss and Shank, 1986). Each test disk was then dampened with 15 μ l of double distilled water and placed in the vial. Control disks dipped into solvent and air dried were similarly treated with double distilled water. Three third instar larvae, starved for 24 hours prior to testing, were placed into each vial. All the feeding tests were conducted for 24 hours in the dark at $65 \pm 2\%$ relative humidity and $29 \pm 2^\circ\text{C}$. Test and control disks were weighed several times to a constant weight before the assay on a Mettler AT261 Delta range balance to ± 0.001 mg. Each test was replicated ten times.

The relative feeding response, X_t/X_o , was calculated for each dose by dividing the mean weight of the treated disks consumed (X_t) by that of the control disks consumed (X_o).

(b) ***Chilo partellus* oviposition bioassay;** These were conducted in a circular chamber, (Kumar and Saxena, 1985; and Ramachandran and Saxena, 1991). The extract or compound was impregnated on a semi circular Piece of filter paper (Whitman Filter paper No. 1, 9cm. diameter) covering half the area, a similar paper without extract covering the other half and serving as the control. A single ovipositing female was released within the circular

chamber overnight and the number of egg masses and eggs laid on the treated and the control semi circles were counted. The percentage preference / non-preference was calculated by dividing the difference in the number of eggs laid on the treated and the control by the total number of eggs laid, $(T-B/T+B)100$, The percentage number of insects laying one and a half (1.5) times or more as many eggs on the blank/treated as on the treated/blank was also calculated. Each test dose was replicated 15-20 times.

All the oviposition tests were conducted for 12 hours in the dark (at night) at $60 \pm 5\%$ relative humidity and $29 \pm 2^{\circ}\text{C}$.

3.3 Plant material:

Whorls of maize; cultivars MP 704 (Resistant) and INBRED A (susceptible), and sorghum; cultivars IS 1044 (resistant) IS 18520 (tolerant) and IS 18363 (susceptible) were obtained from the screen house at the Mbita Point Field Station.

3.4 Insects:


Freshly molted third instar larvae and freshly emerging pupae, of *Chilo partellus*, were obtained from stock cultures maintained on an artificial diet at the I.C.I.P.E insectary (Ochieng *et al.* 1985). Third instar larvae feed first on the leaves of the host plant before gradually boring into the stem. This stage was selected for bioassay owing to the long-term objective of elucidating the basis for the change in the larval feeding sites. The freshly emerged pupae were sexed and paired (male and female) for mating the first night after emerging and oviposition assayed the first night after mating (Kumar and Saxena, 1985).

3.5 Extraction:

The plants were extracted three weeks after emergence. The whorl leaves of each of the maize and sorghum cultivars were extracted by successively soaking them in petroleum ether (20min), ethyl acetate (20 min) and methanol, 20min (for sorghum / oviposition work) and 24 hours (for maize / feeding work). The extract solutions of pet-ether and ethyl acetate were filtered and dried with anhydrous sodium sulphate then concentrated to dryness *in vacuo* at low temperature (40°C). The methanol extracts were filtered and concentrated to a small volume and freeze dried to remove residual water. The residual plant material was also freeze dried to obtain the weight of the dry plant material.

3.6 Fractionation and isolation:

The summary of the fractionation of the extracts and isolation of the constituent allelochemicals is outlined in the diagram below. Fresh maize and sorghum whorls were sequentially extracted in petroleum ether, ethyl acetate and methanol. The extracts were then subjected to chromatographic techniques to isolate the relevant allelochemicals.



3.6.1 Fractionation and isolation of IS 1044 pet-ether extract compounds:

The IS 1044 petroleum ether extract was subjected to column chromatography yielding three main fractions, (I, II & III). Fraction I was further analyzed by GC-MS and by co-injection with authentic standards in the GC.

Fraction III was found to be predominantly one compound by GC analysis and was further purified by column chromatography using a smaller column to obtain the pure compound for Mass Spectroscopic, Nuclear Magnetic Resonance and Infra red analysis.

3.6.2 Isolation and purification of MP 704 ethyl acetate extract compounds:

The allelochemicals of the MP 704 ethyl acetate extract were isolated and purified by preparative High Performance Liquid Chromatography (HPLC). The ethyl acetate extract (each of 100mg) was dissolved in 5ml of methanol and filtered. The sample (50 μ l) was injected into the HPLC Varian 5000 LC. The column used was MCH-10 reverse phase (50cm x 8mm). The solvent system was 30% methanol-water, the flow rate was 3ml/min and the detector was 240nm. The fractions obtained were reduced to small volumes *in vacuo* and freeze dried to remove residual water and obtain the allelocompounds.

4.0 REFERENCES:

Adams C. M. and Bernays E.A. (1978); The effect of combinations of deterrents on the feeding behaviour of *Locusta Migratoria*. Ent. Exp. and Appl. **23**, pp 101 -109.

Alighali, A.M and K.N. Saxena (1983); Larval movement, feeding and development of *Chilo partellus* (Swinhoe) on two sorghum cultivars. Insect Sci. App. Vol **9**. No 1., pp 7-11

Alighali, A.M. (1985); The spotted stalk-borer, *Chilo partellus* (Swinhoe) (Lepidoptera : Pyralidae) and its principle Host, Sorghum. Insect Sci. Appl. vol. **6** no.3, pp 315-322.

Alighali, A.M (1986); Effects of cultivar, time and amount of *Chilo partellus* (Swine hoe) infestation on sorghum yield components in Kenya. Trop pest management, **32**; 126-129.

Ampofo, J.K.O. (1986); *Chilo partellus* (Swinhoe) oviposition on susceptible and Resistant Maize Genotypes. Ins Sci Appl Vol **6** No 3, pp. 323-330

Ampofo, J.K.O. and Nyangiri, E.O., (1986); Maize resistance to *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae): Behaviour of newly hatched larvae and movement from oviposition sites to feeding sites. Appl. Ent. Zool. vol **21** (2), pp. 269-276.

Atkin, D.J.S. and Hamilton, R.J. (1982); The surface of sorghum, The plant cuticle (ed. by D.F. Cutler, K.L. Alvin and C.E. Price), pp. 231-237. Academic Press, London.

Auclair, J.L.; Maltias, J.B. and J.J. Cartier (1957); Factors in a resistance of a peas to the pea aphid *Acyrtiophon pisuim* (Haw) II Amino acids can Entomol **89.**, pp. 457 - 454

Baker, J. E., Rainay, D.P., Norris, D.M., and Strong, F.M. (1986); *p*-hydroxybenzaldehyde and other related phenolics as feeding stimulants for the smaller European elm bark beetle. For. Sci. vol. **14**, pp 91-95.

Barbosa, and V.A. Krischcha, (1987); Am. Natu. Vol. 130 pp 53. In Natural Product reports 1989, ed. J.B. Harborne. pp. 85-109.

Beevor, P.S., D.R. Hall, B.F. Nesbit, V.A. Dyck, G. Arida, P.C. Lippold and A. Oloumi-Sadeghi, (1977); Field trials of synthetic sex pheromones of the striped rice borer *Chilo Suppressalis* (WALKER) (Lidoptera; Pyralidae) and of related compounds. Bull. Entomol. Res. **67**, pp. 439-449.

Beevor, P.S., David, H. and Jones O.T. (1990); Female sex pheromones of *Chilo SPP*, (Lepidoptera: Pyralidae) and their development in pest control. Insect sci. Appl. Vol. **4/5**, pp. 787-794.

Bernays, E. A., Blanay, W. M., Chapman R. F. and Cook, A. G. (1976); The ability of *Locusta migratoria* to perceive plant surface waxes. Symposia Biologica Hangarica, **16**, pp. 35-40.

Bernays, E.A. and Chapman, R.F. (1977); Deterrent chemicals as a basis of oligophy in *Locusta migratoria*. Ecol. Ent. Vol. 2, pp. 1-18.

Bernays, E.A. Chapman, R.F, Mccafferry, A.R, Modder, W.W.D, and Leather, E.M. (1977); The relationship of *Zonocerus variegatus* (L) with cassava (*Manihot esculenta* cranz). Bull. Ent. Res. 67, pp. 427-437.

Bernays, E.A., R.F. Chapman and S. Woodhead, (1983); Behaviour of newly hatched larvae of *Chilo partellus* (Swinhoe) (Lepidopteran:Pyralidae) associated with their establishment in the host plant, sorghum. Bull. Ent. Res. vol. 73, pp. 75-83.

Bianchi, G. Avato, and Salimini, F., (1975); Glossy mutants of maize (IV) Chemical constituents of Glossy-2 Epicuticular waxes. Maydica xx, pp. 165-173.

Bianchi, G., Avato, P., Bertorelli, P. and Mariani, G. (1978); Epicuticular waxes of two sorghum varieties. Phytochemistry, vol. 17, pp. 999-1001.

Binder, R.G., Chan, B.G. and Elliger, C.A. (1979); Antibiotic effects of C₁₀ - C₁₂ fatty acid esters on pink bollworm hollworm and tobacco budworm. Agricultural and biological chemistry. vol. 43, pp. 2467-2471.

Blanay, W.M. and Chapman, R.F. (1970); The functions of the maxillary palps of Acrididae (Othoptera). Entomologia Experimentalis et Applicata, vol. 13, pp. 363-376.

Broughtton, H.B., S.V. Lay, A.M.Z. Slawin, D.J. Williams and E.D. Morgan, (1986); J. Chem. Soc. Chem. Commun. **46**. In Natural Product reports 1989, ed. J.B. Harborne, pp. 85-109.

Butterworth, J.H, Morgan, E.D. and Percy, G.R. (1972); The structure of azadirachtin, the functional groups. J. Chem. Soc. Perkin Trans I. No. **19**, pp. 2445-2450.

Buttery Ron G. and Lousia, C. Ling. (1984); Corn leaf volatiles. Identification using tenax trapping: J. Agr and Food chem (1984), pp. 1104 - 1106.

Buttery Ron G. and Louisa C. Ling, (1985); volatile components of corn roots . Possible insect attractants. J. Agric. foodchem. July-August (1985), pp. 772-774.

Chandravadana, M.V. (1987) J. Chem. Ecol. vol. **13** pp 1689; In Natural Product reports 1989, ed. J.B. Harborne, pp. 85-109.

Chapman R.F. (1977); The role of leave surface in food selection by acridids and other insects. Colloques Internationaux du C.N.R.S. **226**, pp. 33 - 149.

Cooper Driver, G.A. and Swain, T. (1976); Cynogenic polymorphsim in bracken in relation to herbivore predation. Nature land. **260**, pp. 640.

Derridj, S and Fiala V. (1983); Sucres solubles. des Feuilles de mais (*Zea mayl*) et oviposition de la pyrale (*Ostrinia nubilalis* HBn) C.r. hebd seanc Acad Agric Fr **99**, pp. 465 - 472.

Dabrowski, Z.T. and E.L. Kidiavai,(1983); Resistance of some sorghum line to the spotted stalk borer *chilo partellus* under Western Kenya conditions. Insect sci. Appl. Vol. 4, pp. 119-126.

Dethier, V.G (1947); Chemical insect attractant and repellent. Blakiston Philadelphia.

Dethier, V.G. (1966); Feeding behaviour. In, Insect behaviour (Haskell, P.T., ed.) RES. symposium series No. 3, pp. 46-59.

Dethier, V.G. (1980); Evolution of receptor sensitivity to secondary plant substances with special reference to deterrents. Am. Nat. 115, pp. 45-66.

Dethier, V.B (1982); Mechanisms of host-plant recognition. Entomol Exp Appl 31, pp. 49-56.

Doss, R.P., and Shanks, C.H. (1986); use of membrane filters as substrate in insect feeding bioassays. Bull. Entomol. Soc. Am. 32, pp. 248-249.

Dreyer, D.L., Reese, J.C., and Jones, K.C (1981); Aphid feeding deterrents in sorghum. Bioassay, isolation and characterization. J. chem. Ecol. vol. 7, pp. 273-284.

Eglinton, G. and Hamilton, R.J. (1963); The distribution of alkanes, In Chemical plant taxonomy (ed. by T. Swain), pp. 187-217. Academic Press London.

Feeny, P. (1988); *Phytochemistry* Vol. 27 pp. 3439, In *Natural Product reports 1989*, ed. J.B. Harborne, pp. 85-109.

Fisk, J. (1980); Effects of HCN, phenolic acids and related compounds in sorghum bicolor on the feeding behaviour of the planthopper *Peregrinus maidis*. *Entomol. Exp. Appl.* vol. 27, pp. 211-222.

Gnanasunderam, C. and O.R.W. Sutherland, (1986); *Phytochemistry* vol. 25 pp 409, In *Natural Product reports 1989*, ed. J.B. Harborne, pp. 85-109.

Haskins, F.A and H.J Gorz (1984); Dhurrin and p-hydroxybenzaldehyde in seedlings of various Sorghum species. *Phytochem.* Vol. 24, pp. 597-598.

Hedins, P.A. Jenkins, J.N. and Maxwell, F.G. (1977); Behavioral and development factors affecting host plant resistance to insects. In *Host plant resistance to pests*, (Hedin, P.A. ed.) pp. 231- 275. Washington .D.C.

Hill, D.S. (1983); *Agricultural insect pest of tropics and their control*. University Press Cambrigde pp. 57.

Huesing and D.Jones, (1987); *Phytochemistry* vol.26 pp. 1381. In *Natural Product reports 1989*, ed. J.B. Harborne, pp. 85-109.

I.C.I.P.E annual report (1988); pp. 11.

Janzen, D.H (1979); New Horizons in the biology of plant defense pp. 331-350.

Jones, K.C. and Klocke, D.A, (1987); Aphid feeding deterency of ellagitannin; their phenolic hydrolysis products and related phenolic derivatives. Ent. Exp. **44**, pp. 229-234.

Jotwani, M.G. (1981); Insect resistance in sorghum plants. Insect Sci Appl. **2**, pp. 93-98.

Jotwani, M.G, S. Chaudhary and S.P.Singh, (1978); Mechanisms of resistance to *Chilo partellus* (swinhoe) in sorghum. Indian J. Ent. **40**, pp. 273-276.

Jotwani, M.G. & J. C. Davies, (1980); Insect resistance studies on sorghum at International institutes and National programs with special reference to India. In biology and breeding for resistance to anthropoids and pathogens in Agricultural plants. Ed. Harris, M.K. Texas, A and M. Univ. systems college station, Taxes, pp. 224-226.

Kalode, M.B and Pant, N.C. (1966); Studies on the suitability of different varieties of sorghum, maize, Bajra to *Chilo Zenellus* (Swinhoe) under field and cage conditions and the methods of determining it. Ind. J Ent. **28**, pp. 458 - 464.

Kira, M.T ; Guthrie, W.D. and Huggans, J.L. (1968); Effects of drinking water on production of eggs by the European Corn bore .J Econ. Ent . **62**, pp. 1366 - 1368.

Klingauf, von F., Nocker-Wenzel, K. and Klain, W. (1971); Einfluss einigers Wachskomponenten von *Vicia faba* auf das Wirtswahlverhalten von *Acyrtosiphon pisum*

(Harris) (homoptera: Aphididae). Zeitschrift fur Pflanzenkheiten und Pflanzenschutz, **78**, pp. 641-648.

Kogen, M. and Ortman. E.F. (1978); Antixenosis, A New term proposed to replace painter's non - preference modality of resistance . ESA Bull Vol **24**.

Kumar, H. and K.N Saxena (1985); Ovipositional response of *Chilo partellus* (Swinhoe) to certain susceptible and resistant maize genotypes. Insect Sci. Appl. Vol **6** (3), pp. 331-335.

Kumar, H. & Saxena, K.N. (1985); Oviposition by *Chilo partellus* (swinhoe) in relation to its mating diurnal cycle and certain non-plant surfaces. Appl Ent. Zool. **20**, PP. 218 - 221.

Kumar, H. (1986); Enhancement of ovipositional responses of the stem-borer *Chilo partellus* (swim hoe) to certain Sorghum cultivars in relation to their resistance or susceptibility insects - plants, PP. 313 -318.

Lal, G. and J.C. Pant, (1980); Laboratory and field testing for resistance in maize and sorghum varieties to *Chilo partellus* (swinhoe). Indian J. Ent. Vol. **42**, pp. 606-610.

Lwande, W. and M.D. Bentley.(1987); Volatiles of Sorghum Bicolor seedlings J.Natural products Vol **50** NK5. pp. 950 - 952.

Maltais, J.B. (1951); Nitrogen content of different varieties of peas as a factor affecting infestation by *Macrosiphum pisi* (Kith) a preliminary report. *Can. Entomol.* **83**, pp. 29.

Martin, J.T. and Juniper, B.E. (1970); *The cuticles of plants*. Edward Arnold, London.

Matsuda, K. and Senbo. (1986); Chlorogenic acid as a feeding deterrent for salicaceae feeding leaf beetle. *Apple Entomol Zool* **21**, pp. 411-416.

Maxwell, F.G. Schuster, M.F; Meredith, W.R. and M.L. Laster (1976); Influence of the Nectariless character in cotton on harmful and beneficial insects. pp. 157-161.

Miller, J.R. and Sticker, K.L. (1984); Finding and accepting host plants in chemical Ecology of insects. pp. 128-158.

Munakata, A.K. (1977); Insect antifeedants of *Spodoptera litura* in plants. In, *Host plant resistance to pests*. (Hedin, P.A. ed.), pp. 185-196. Washington D.C.

Neimeyer, H.M, Pesel, E., Copaya, S.V.Bravo, R., Franko, S. and Franko, W. (1989); Changes in hydroxyamic acid levels of wheat plant induced by Aphid feeding. *Phytochem.* Vol. **28**, pp. 447-449.

Norris, D.M (1986), Antifeeding compounds. In, *Sterol biosynthesis, Inhibitors and Antifeeding compounds*. (Kato, T., Kramer, W., Kuck, H.H., Norris D.M. and Scheinpflug, H., contributors), pp. 96-246.

Norris, D.M. and Kogen, M.(1980); Biochemical and morphological bases of resistance in breeding plant resistant to insects. pp. 23-61.

Ochieng, R.S., Onyango, F.O, and Bungu, M.D.O, (1985); Improvement of techniques for mass culture of *Chilo partellus* Swinhoe). Insect sci. Appl. Vol. 6, pp. 425-428.

Odindo, M.O, (1991); Management of cereal stem borers, especially *Chilo partellus*, using microsporidia. Insect sci. Appl. Vol. 2, No. 1/2/3, pp. 51-55.

Painter, R.H. (1958); Resistance of plants to insects. A Rev Ent.3, pp. 267-290.

Painter, R.H. (1951); Insect resistance in crop plants. University press of Kensas Lawrence Kensas.

Pathak, M.D. and Dale, A.A. (1983); The biochemical basis of resistance in host plants to insect pests. The New Frontier CHEMRAWN II. Ed dy Schemilt L.W, pp. 129-142.

Pathak, R.S. (1991); Genetic expression of the spotted stem borer *Chilo partellus* (swinhoe) Resistance in three maize crosses. Insect sci. Appl. Vol. 12, No. 1/2/3, pp. 147-151.

Quraishi, M.S. (1971); Toxic and teratogenic effects of saturated and unsaturated fatty acid on house fly larvae. Journal of Economic Entomology, 64, pp. 787-792.

Raina Ashok, K. (1981); Deterrence of repeated oviposition in sorghum shootfly (*Atherigona soccata*). J. Chem. Ecol. vol. 7, No. 5, pp. 785-790.

Raina Ashok, K. (1982); Fecundity and oviposition behaviour of the sorghum shootfly (*A. soccata*). Entomol. Exp. & Appl. Vol. 31, pp. 381-385.

Ramachandran, R. and K.N. Saxena. (1991); Some aspects of emergence, mating, oviposition and flight activity of *Chilo partellus* swinhoe (Lep. Pyralidae) J. Appl Ent. 112, pp. 43 - 52.

Ramamurthi Jambunathan, Larry, G. Butter, Ranajit Bandyopdhyay and Lewis , K. Mughogho, (1986) Polyphenol concentrations in grain, leaf and callus tissues of mold susceptible and mold resistant sorghum cultivars. L. Agri. foodchem. Vol. 34, pp. 425-429.

Rene Gujer, Daniele Magnolato and Rone Salf, (1986); Glucosylated flavonoids and their phenolic compounds from sorghum. Phytochem. Vol. 25 No. 6, pp. 1431-1436.

Robinson, J.F., J.A. Klun and T.A. Brindley (1978); European corn borer: A no preference mechanisms of feeding resistance and its relationship to 1,4-Benzoxazin-3-one concentration indent corn tissue. J.Econ. Entomol. vol. 71, pp. 461-465.

Saxena, K.N. (1969); Patterns of insect plant relationships determining susceptibility/resistance of different plants to an insect . entomol. Exp. & Appl. Vol. 12, pp. 751-766.

Saxena, K.N. and Thorsteinson, A.J. (1971); Effects of synthetic "Queen substance" and analogues on survival, molting and metamorphosis of *aedes aegypti*. J. Econ. Entom. **64**, pp. 287-291.

Saxena, K.N. (1985); Behavioral Basis of plant resistance or susceptibility to insects. Insc Sci App Vol **6**, No 3, pp. 303-313.

Saxena, K.N. (1987); Ovipositional responses of the stem-borer *Chilo partellus* (Swinhoe) to certain sorghum cultivars in relation to their resistance or susceptibility. In insects - Plants. Eds. Labeyrie, Fabres, Lachaise and Junk; Publ. Dordrecht, The Netherlands. pp. 313-318.

Saxena, K.N. (1990); Mechanisms of resistance/susceptibility of certain sorghum cultivars to the stem borer *Chilo partellus*; role of behaviour and development. Entomol. exp. & appl. vol **55**, pp. 91-99.

Schoonhoven, L.M. (1972); Secondary plant substances and insects. Rec. Adv. Phytochem. Vol. **5**, pp. 197-224.

Schoonhoven, L.M. and Jermy, T. (1977); A behavioral and electrophysiological analysis of insect feeding deterrents. In Crop protection agents; Their Biological evolution. (Macfarlane, N.R, ed), pp. 133-146. Academic press New York.

Schoonhoven, L.M. (1982); Biological aspects of antifeedants. Entomol. Exp. & Appl. Vol. **31**, pp. 57-69.

Seshu Reddy, K.V. (1982); Pest management in sorghum (II) proceedings of the international symp. on sorghum 12-7 Nov 1981, pp. 238 - 245.

Seshu Reddy K.V. (1985); Relative susceptibility and resistance of some sorghum lines to stem-bores in western Kenya. *Ins Appl* **6**, pp. 401-404.

Seshu,R.K.V. (1988); An assessment of on-farm yield losses in sorghum due to insect pests. *Insect Sci. Appl.* vol.9 no.6, pp. 679-685.

Seshu Reddy, K.V and K.O.S.Sum, (1991); Determination of economic injury level of the stem borer *Chilo partellus* (swinhoe) in maize *Zea mays* L. *Insect sci. Appl.* Vol. **12** No. 1/2/3, pp. 269-274.

Seshu Reddy, K.V, (1991); Insect pests of sorghum in Africa. *Insect sci. Appl.* Vol. **12** No. 5/6, pp. 653-657.

Silverstein, R.M. and J.B. Simeone (eds) (1983); North American symposium on Allelopathy *J. chem Ecol.* **9**, pp. 935 - 1292.

Silverstein, R.M.; G. Clayton Bassler and Torence C. Morrill (1981); Spectrometric identification of organic compounds. Fourth Edition. John Willey & Sons Newyork. Chichester. Bristane. Toronto. Singapore. pp. 3-278.

Singh, S.R. (1978); Plant resistance and pest management. ICIPE /UNEP Group training course, Lecture Notes for 1978 course 16th July - 6th August.

Singh, B.U, B.S. Raina, B.B. Reddy and N.G.P. Rao (1983); Host plant resistance to stalk borer *Chilo partellus* (swin). In sorghum; Insect sci. Applic. Vol. 4, pp. 407-413.

Singh, B.U. and Raina, B.S. (1989); Varietal resistance in sorghum to spotted stem borer *C.partellus* (swinhoe). Ins Sci App. 10, pp. 3-27.

Smith, C.M; Wilson, R.F. and Brim, C.A. (1979); Feeding behaviour of Mexican bean beetle on leaf extracts of resistant and susceptible soybean genotypes. J.econ. Ent. 72, pp. 374-377.

Sogawa, K. and Pathak, M.D. (1970); Mechanisms of brown planthopper resistance to Mudgo variety of rice (*Hemiptera Delphaudae*) Appl Ent. Zool 5, pp. 145-158.

Stadler, E. (1978); Chemoreception of host plant chemicals by ovipositing females of *Delia (Hylemya brassicae)*. Entomologia experimentalis et Applicata. 24, pp. 711-720.

Stadler, E. (1982); Sensory physiology of insect plant relationship -round - table Discussion. In; Proc. - 5th Int. sym. Insect-plant relationships. (Visser, J.H. and Minks, A.K, eds.), pp. 81-91. Pudoc, Wageningen.

Stockel, J. M; Bar, J. N. Boidron and G. Bourgeois (1987); Methodological approach to identify chemical oviposition stimulants from maize for European corn borer. J. chem. Ecol. vol 13 (3), pp. 557 - 564.

Teetes, G.L. Seshu Reddy, K.V. , Leuschner, K.and House, L.R. (1983); International Crop Research Institute for the Semi-Arid Tropic; info. Bul. Vol. 12, pp. 46-49. Patancheru. A.P. India.

Thorstein, A.J. (1960); Host selection in phytophagous insects A Rev. Ent. 5, pp. 193-218.

Torto, B., Hassanali, A. and Kailash, N.S. (1990); Chemical aspects of *Chilo partellus* feeding on certain sorghum cultivars Ins App Vol 11 . No 4/5, pp. 649 - 655.

Torto, B., A.Hassanali; Kailash N.S and Sagary Nokoe (1991); feeding responses of *C. partellus* (Swine hoe) larvae to sorghum plant phenolics and their analogs. Journal of chemical Ecology Vol 17 No 1, pp. 66-78.

Unnithan, G.C; K.N. Saxena; M.D. Bentley and A. Hassanali (1987); Role of sorghum extracts in Eliciting oviposition on 2 non-host by the Sorghum shootfly Env. Ento. 16. pp. 967 - 970.

Unnithan, G.C. and K.N. Saxena. (1990); Diversion of oviposition by *Atherigona Socata* to Non - host maize with Sorghum Seedlings extract Env. Entomol. 19 (5), pp. 1432 - 1437.

Unnithan, G.C. and K.N. Saxena (1991); Pheromonal trapping of *Chilo partellus* (swinhoe) (Lepidoptera: Pyralidae) Moths in relation to male population density and competition with females. Appl. Ent. Zool. **26** no.1, pp. 17-28.

Van Hamburg H. (1980); The grain Sorghum stalk borer *Chilo partellus*. Survival and location of Larvae at different infestation levels in plants of different ages. J. Ent. Soc sth. Afr. **43**; pp. 71-76.

Waladde, S.M ; S.A. Ochieng and H.M. Kahoro (1989); Responses of *Chilo partellus* Larvae to host plant materials: ins. sci. Appl. **11** (1), pp. 79 - 89.

Wieczorek H. (1976); The glucoside receptor of the Larvae of *Mamestra brassicae* L. (Lepidoptera Noctuidae) J. Comp. Physiol. (A) pp. 106 - 176.

Wilkinson, R.E. and Cummins,D.G. (1981); Epicuticular fatty acid, fatty alcohol and alkane contents of bloom and bloomless sorghum "redbine 60" leaves. Crop science, **21**, pp. 397-399.

Whitman, D.W. (1988); Allelochemical Interactions among plants, herbivores and their predators. pp. 11-64.

Woodhead, S. and Bernays, E.A. (1978); The chemical bases of resistance of sorghum bicolor to attack by *L. migratoria*. Ent. Exp & Appl. Vol. **24**, pp. 123-144.

Woodhead, S. (1982); p-Hydroxybenzaldehyde in the surface wax of sorghum: its importance in seedling resistance to acridids. *Entomologia Experimentalis et Applicata*, **31**, pp. 397-399.

Woodhead, S. (1983); Surface chemistry of sorghum Bicolor and its importance in feeding by *L. Migratoria*, *physiol, Entomol.* Vol **8**, pp. 345 - 352.

Woodhead, S. and Janeja, S.K. (1987); The importance of the behaviour of young larvae in Sorghum resistance to *Chilo partellus*. *Entimologia Experimentalis et. appl.* **45**, pp. 43-54.

Woodhead, S. and D.E. Padgham, (1988); The effect of plant surface characteristics on resistance of rice to brown planthopper (*Nilaparvta lugens*). *Entomol. Exp. & Appl.* Vol. **44**, pp. 15-22.

Young W.R and Teetes G.L (1974); Sorghum entomology: A Rev. *Ent.* **22**, pp. 193-219.