

*icipe* Tropical Insect Science for Development

**A**  
**REPORT OF THE**  
**DELIBERATIONS OF THE**  
**MALARIA TASK FORCE**  
**MEETING**

*Identifying Research Priorities in Vector Biology and Ecology for  
Integrated Malaria Control in Africa*



ICIPE HEADQUARTERS,  
DUDUVILLE, KASARANI, NAIROBI

4-8TH FEBRUARY 1996

*icipe* Tropical Insect Science for Development

**A  
REPORT OF THE  
DELIBERATIONS OF THE  
MALARIA TASK FORCE  
MEETING**

*Identifying Research Priorities in Vector Biology and Ecology for  
Integrated Malaria Control in Africa*



ICIPE HEADQUARTERS,  
DUDUVILLE, KASARANI, NAIROBI

4-8TH FEBRUARY 1996



## EXECUTIVE SUMMARY

Of mosquito-borne diseases — malaria, filariasis, dengue, yellow fever and Japanese encephalitis — malaria is by far the most important in terms of the number of individuals it affects and the mortality it causes. It affects nearly half the world population (2.2 billion people) in some 90 countries, with 300-500 million estimated clinical cases each year, 1.5-3.0 million deaths, 90% of which occur in tropical Africa. The worsening malaria situation in the world led the WHO to declare malaria control of global priority. However, incidences of other diseases have also been on the increase in some areas. Vector control is acknowledged as an essential part of control programmes of these diseases to which ICIPE can make vital contributions.

In February 1996, ICIPE together with WHO and the Netherlands government sponsored a Task Force Meeting of mosquito and malaria experts from Africa, USA, Europe and Asia to review the status of malaria vector research and to identify critical gaps which ICIPE could address. A unanimous conclusion of the meeting was that much of the information needed for effective vector control focussed on disease control is incomplete, and, in some important aspects, lacking. Surprisingly little is known about the relation between inter and intra-specific diversity among mosquito species, their vectorial dynamics, and underlying population genetics, nor between disease incidence and population dynamics of critical mosquito populations in different environments. A recurring paradox in Africa is apparent lack of direct relation between transmission pressure and malaria disease. Little is known about larval population ecology and physiology, and particularly the relation between environment, competitive population dynamics of different sibling species (e.g. of *Anopheles gambiae*) and vectorial fitness of adults. As a result, no meaningful ecosystem based quantitative models can be formulated to guide proactive vector control strategies. Moreover, some of the key behaviours of malaria vectors, such as host location and oviposition, and their mediating cues, have not been investigated in any detail. The discovery of semiochemical cues from such research could provide powerful tools for behavioural manipulation of critical vectors.

## NEEDS AND KNOWLEDGE GAPS

### *Gaps in malaria vector biology and ecology*

1. Among the factors which make the control of malaria in Africa especially imperative and challenging are the rising resistance to drugs and pesticides; outbreaks of the disease in areas such as the Kenya highlands that were formerly malaria-free; influence of abiotic factors such as climatic change, civil strife and urban migration on disease incidence;



theirs? What attracts them to each other to form the mating pair? How are breeding sites found and selected? What is the role of visual and olfactory (chemical) cues in this respect?

- (vi) How do the adults locate their hosts (host-finding behaviour)? What chemical cues or other factors emanating from the host or the environment are involved?
- (vii) Is there a difference in the behaviour of resistant and non-resistant mosquitoes?
- (viii) What are the physiological effects of attractants and repellents? Can control methods such as bednets, for instance, produce an avoidance response?
- (ix) What is the effect of parasites on the adult mosquito?

5. There is a gap in knowledge about species **distribution and stratification**. There have been several admirable efforts at mapping species, such as that by the South African Institute for Medical Research, and stratification based on endemicity levels and vector species done by WHO in South Africa, Mauritius, Somalia and Sudan, but this is needed in other countries as well.

6. Although there is some basic knowledge about the systematics and biology and ecology of the terrestrial adult anophelines, very little is known about their **aquatic larval population ecology and physiology**. Although research in this area is admittedly difficult, questions that need to be answered include the following:

- (i) What is the role of mosquito larvae in the food web and ecology of water-based systems?
- (ii) What is the relative contribution of different feeding sites to overall disease transmission pressure?
- (iii) What is the basis of the competition between the *Anopheles gambiae* sibling species?
- (iv) What is the nature and role of larval feeding behaviour in population dynamics? Is the vectorial fitness of the adult related to larval size?

7. Almost no biological information exists about the **egg stage** of the mosquito, yet this could offer an avenue for possible vector control. This is an obvious gap in knowledge.

8. Although laudable work on vector identification has been done in Africa utilising a variety of techniques including PCR (polymerase chain reaction) and DNA probes, the **lack of a reliable vector species identification service** in Africa constitutes a major research handicap. There is an urgent

need for a vector species identification centre. In order to be effective, such an identification centre should have well equipped laboratories, reliable services, trained staff, an insectary, a mosquito reference collection, and a well equipped library.

9. There is a need for development and **improvement of the PCR** technique for species identification and for the development of molecular biology techniques for the rapid detection of mosquito resistance to pesticides and of the parasite to chemotherapy.
10. The nature of **vector-parasite interactions**, especially at the molecular level, and the mechanisms underlying these interactions, is not well understood. Knowledge of these interactions could reveal vulnerable points which could be manipulated to disrupt key biological processes that allow parasite development and transmission. Some of the questions to be answered include the following:
  - (i) What is the role of trypsin in parasite development?
  - (ii) Does the inhibition of ookininetes binding to lectin-based molecules open the way for “glycotherapy” as a method of control?
  - (iii) Can transmission-blocking antibodies be developed and used for control?
  - (iv) What is the cell biology of the infection process, and how does this result in vector specificity?
11. The relationship between **transmission pressure and malaria disease** is not well understood. Recent epidemiological studies confirm that malaria vector competence varies across species. But, whereas it has always been assumed that there is a direct relationship between the intensity of malaria (malaria rate/incidence) and the entomological inoculation rate (EIR), recent studies in Kenya suggest that the malaria rate is not necessarily a function of EIR. Studies on the transmission rate and the incidence of severe malaria at the Kenya Coast found that although the biting rates were very low, the incidence of severe malaria was unusually high, and exposure to even one infected mosquito can result in severe malaria. The situation for Kisumu in western Kenya was the opposite: despite a very high EIR, cases of severe malaria were comparatively few. There is need for further studies to shed more light on this apparent paradox.
12. Current vector control methods rely largely on synthetic insecticides or permethrin/deltamethrin-impregnated bednets. Both these methods have failed to effectively interrupt malaria transmission, especially in

holoendemic areas. Furthermore, eradication or complete interruption of malaria transmission is neither feasible nor desirable in holoendemic areas, as it will allow for the maintenance of natural immunity. Conventional methods of control need to be re-examined in light of the foregoing. Some questions that remain to be answered include the following:

- (i) In the case of bednets, do they really reduce deaths in areas of high transmission?
- (ii) Will the most vulnerable groups, pregnant women and children become more susceptible to malaria if they stop using bednets?
- (iii) What are more effective methods of fabric impregnation?
- (iv) Is the use of bednets producing mosquito tolerance to pyrethroids, as reports from Kenya and Dubai suggest?

13. The use of non-conventional control or repellent substances such as neem and other botanicals need to be investigated, in light of their traditional uses in some cases and their generally lower costs and local availability.

14. Applied research without sound theoretical underpinnings is not sustainable. There is need to carefully balance strategic and applied issues in the design of mosquito and malaria control projects.

#### *Needs for capacity building and collaboration in malaria vector research*

15. Training in all aspects of insect science, with a specialisation on malaria vectors is needed at all levels: for postgraduate(MSc and PhD) students, postdoctorate fellows, technicians, and management and supervisory staff. Postdoctorate trainees should be attached to on-going projects.

16. It is important for African training institutions to link up with each other and with institutions in the North and in other countries of the South, for maximum sharing of resources and expertise. This can be achieved, for instance, by exchange of staff, research students, postdoctorates and visiting scientists. This is already being done in South Africa and at the ICIPE through the African Regional Postgraduate Programme in Insect Science (ARPPIS). ARPPIS is a training network of about 30 universities in Africa who collaborate with ICIPE in training students at PhD and MSc.levels in insect science.

17. Links between researchers of international organisations and those of national organisations have historically been weak or even non-existent in most of Africa. There is need to establish and strengthen collaboration between the two groups.



18. African institutions suffer from a plethora of problems, including poor supervision due to the high student to staff ratio; poor general facilities, utilities and infrastructure; lack of specialised research facilities such as an insectory and reference collections; and a general lack of funds for conducting research. There is a need to **identify and equip "centres of excellence"**, which would serve national programmes throughout Africa, and act as incubation points for the development of national research capabilities.
19. There is need for a **multidisciplinary approach** to training and capacity building, with social scientists and epidemiologists, for instance, working together with entomologists. An integrated approach that brings together agricultural, medical and forest entomology will be required in order to fully address the problem of malaria in Africa.

## **RECOMMENDATIONS FOR ICIPE'S PROGRAMME DEVELOPMENT**

20. The Task Force Meeting participants unanimously endorsed ICIPE's role in malaria vector research, in four general activity areas. The research orientation should be problem solving, innovative and aimed at filling in the knowledge gaps. It should also be multidisciplinary, network-coordinated research with a well-defined bandwidth. The four areas are:
  - (i) basic strategic research whose output is relevant to malaria control in Africa
  - (ii) operational research, through networking with African national institutions
  - (iii) research capacity building
  - (iv) provision of services, such as species identification, resistance monitoring, etc.
21. The over-riding strategy recommended by the participants was that ICIPE should develop *novel* concepts for eventual use in control, i.e., no bednets, no insecticides, and no drugs.

### ***Recommendations for strategic research in integrated vector and disease control***

22. All activities and projects undertaken relating to strategic research should include the following principles:
  - (i) Incorporation of social and economic considerations in the development, implementation, and assessment of control

strategies. ICIPE's Social Science Department would play a major role here and in (ii) & (iii) below.

- (ii) Stratification of malaria eco-epidemiological zones in preparation for control operations.
- (iii) Assessment of the effectiveness and impact of interventions.

23. ICIPE should undertake basic research with an applied outcome, and with a focus on product development. The products should be tested in the field and in various ecological settings. The scientific direction and product development should be built on current ICIPE interests and upon the needs of the local communities and other target groups.

24. Malaria vector ecology is the basic foundation that provides the theoretical underpinning upon which other vector research components will be built.

The main research components that would together contribute to the development of an integrated mosquito/malaria control programme at ICIPE were identified as follows:

- (i) vector ecology and eco-epidemiology, including effects of environmental change
- (ii) population genetics
- (iii) Behaviour and chemical ecology of both adult and larval stages, especially use of botanicals, odour baits, oviposition pheromones, repellents and attractants.
- (iv) modelling of vector populations and disease dynamics

25. The priority areas for ICIPE's research in genetics and molecular biology research were suggested as being in:

- (i) the genetics and biochemistry of insecticide resistance
- (ii) vector-parasite interactions (in collaboration with other on-going programmes)
- (iii) provision of field support for vaccine development

26. Research in population genetics is important in understanding the mosquito in its natural biological community. The knowledge so gleaned will be valuable to many control strategies.

27. ICIPE's experience in biological control should be utilised by undertaking research into the development of bacteria such as *Bacillus thuringiensis*(Bt) and fungi for vector control.

28. The following specific topics for strategic research needed for effective vector management were recommended:
- (i) Eco-epidemiological zoning, employing GIS and other techniques
  - (ii) Larval ecology, especially the place of *Anopheles* larvae in the food chain, encompassing research on food webs in specific agro-ecosystems
  - (iii) Behavioural biology so as to gain a better understanding of behavioural change, such as for example vector behaviour in the dry season and its impact on transmission dynamics and species competition

### ***Intervention or Operational Research***

29. In the area of operational research, the following topics and activities were recommended:
- (i) development of attractive devices for sampling and control
  - (ii) assessment of the contribution of different vector groups to malaria transmission
  - (iii) monitoring of climate change-induced transmission patterns in highland areas
  - (iv) studies on the man-made micro-environmental change and its impact on mosquitoes and transmission
  - (v) exploration of habitat management as a control strategy
  - (vi) testing the efficacy of insecticides in experimental huts
  - (vii) development of techniques to track the spread of insecticide resistance and especially monitoring of resistance to pyrethroids and cross resistance between pyrethroids
  - (viii) development of genetic control tools that would directly reduce vector fitness or block pathogen transmission.

### ***Capacity Building and Collaboration***

30. The priorities in capacity building and levels of training linked to ICIPE's programme goals were identified as follows:
- (i) PhD - level training, particularly in the research component, through the existing ARPPIS programme
  - (ii) training of middle-level (MSc) workers, particularly in the research component
  - (iii) postdoctoral level training, integrated into on-going malaria research projects

- (iv) specialised short courses

The meeting did not reach a consensus about the need for offering coursework or the training of technicians by ICIPE.

- 31. Networking and collaboration should be undertaken by ICIPE with the following:
  - (i) national research institutes in Africa
  - (ii) CGIAR institutes
  - (iii) the ORSTOM network in Francophone West Africa
  - (iv) international organisations and programmes, such as WHO, TDR, PEEM, etc.
  - (iv) "centres of excellence" in industrialised countries
  - (v) NGO's, such as AMREF

It was recommended that ICIPE collaborate with national research systems, such as the Kenya Medical Research Institute(KEMRI) in this research. Kenya was thought to be a good starting point for collaboration with national institutes because it has a relatively strong national governmental malaria control strategy.

- 33. Mobilisation of funds could be approached through:
  - (i) Policy frameworks such as those of WHO; Agenda 21; the Biodiversity Convention, the Ramsah Convention, etc.
  - (ii) Assessment of non-conventional desks in bilateral agencies
  - (iii) "Packaging" of proposals in line with current priorities of national and international malaria control strategies and funding agencies.

## **ACKNOWLEDGEMENTS**

ICIPE is grateful for the encouragement and support of the Netherlands Government, the Government of Denmark, WHO, and ORSTOM in the Centre's endeavors to re-establish its medical vector research activities.



# PROGRAMME

## A REPORT OF THE DELIBERATIONS OF THE MALARIA TASK FORCE MEETING

*Identifying Research Priorities in Vector Biology and Ecology for  
Integrated Malaria Control in Africa*



**TASK FORCE MEETING TO IDENTIFY AREAS OF RESEARCH IN VECTOR  
BIOLOGY AND ECOLOGY IN SUPPORT OF MALARIA CONTROL IN  
AFRICA**

**Nairobi, 4–8 February 1996  
Venue: Rachel Galun Hall**

***Objectives:***

To review potential methods for the control of malaria vectors, with particular reference to integrated control approaches;

To make recommendations for strategic research activities on fundamental and applied aspects of malaria vector control;

To recommend a suitable institutional framework for effective mosquito research and for the development of integrated control strategies, and to assess the suitability of the ICIPE as the centre for such studies in Africa;

To make recommendations for mobilizing and/or training of suitable personnel, to sustain research and application work on malaria vector control in Africa.

***Task Force Chairmen: Robert Bos and Peter Esbjerg***

**Programme**

**Sunday, 4th February: Arrival of participants**

**Monday, 5th February: Current situation, Research Status and Needs**

- 0800 : Registration  
0830 : Welcome and Opening remarks: **Hans R. Herren, Director General, ICIPE**  
0840 : Malaria in Africa: Current situation and outlook for control (**Robert Bos**)  
0920 : Vector research in aid of malaria control: historical solutions and future developments (**Caroline Maxwell**)  
1000 : COFFEE/TEA



- 1030 : Malaria vector research in Eastern and Southern Africa: laboratory studies (*R. Hunt*)
- 1110 : Malaria vector research in East Africa: field studies (*F. W. Mosha*)
- 1150 : Malaria vector research in West Africa (*Sekou T. Traore*)
- 1230 : Discussion
- 1300 : LUNCH
- 1400 : Need for research in vector biology and ecology for malaria control (*W.A. Hawley*)
- 1430 : ICIPE as International Centre For Vector Research (*Peter Esbjerg*)
- 1500 : Discussion
- 1530 : COFFEE/TEA
- 1610 : First workshop: Needs and constraints in current research in ecology, epidemiology, behaviour, genetics, molecular biology, control; (participants are divided in small groups to discuss different topics as indicated by the Chairmen).
- 1800 : Adjourn

**Tuesday, 6th February : Recent advances: Background**

- 0830 : Recent advances in mosquito genetics (*F. H. Collins*)
- 0900 : Recent advances in mosquito/parasite interactions (*P.F. Billingsley*)
- 0930 : Recent advances in mosquito behaviour (*W. Takken*)
- 1000 : COFFEE/TEA
- 1030 : Recent advances in malaria epidemiology (*J.C. Beier*)
- 1100 : Recent advances in vector control (*S. Lindsay*)
- 1130 : Recent advances in malaria control (*S.K. Subbarao*)
- 1200 : Recent advances on malaria research in Orstom (*D. Fontenille*)
- 1230 : General Discussion
- 1300 : LUNCH
- 1400 : Second workshop: research in integrated vector and disease control in Africa
- 1700 : Adjourn
- 1930 : Dinner to be hosted by ICIPE Director General (Nyama Choma Ranch, Safari Park Hotel)

BLACK

## LIST OF PARTICIPANTS

Dr Amha Aseffa  
Professor and Scientist  
College of Medicine  
Department of Biochemistry  
& Molecular Biology  
Howard University  
Washington, D.C. 20059  
U.S.A.  
Tel: no. 202 806 7606  
Fax: 202 806 5721

Dr E.J. Asimeng  
Medical Entomologist & Parasitologist  
Medipest Consult and Research Ltd.  
Vector Control Specialists  
3 Riverside Drive  
P.O. Box 14756  
Nairobi, Kenya  
Tel: 254-2-441699

Dr John C. Beier  
Associate Professor  
Department of Tropical Medicine  
Tulane University  
1501 Canal Street, Rm. 505  
New Orleans, LA 70112  
U.S.A.  
Tel: 1-504 585 6949  
Fax: 1-504 587 7113  
e-mail: jbeier@mailhost..tcs.tulane.edu

Dr. Peter F. Billingsley  
Department of Zoology  
University of Aberdeen  
Tillydrone Avenue  
Abardeen AB9-2TN  
Scotland  
U.K.  
Tel: 44 1 224 272 882  
Fax: 44 1 224 272 396  
e-mail: p.billingsley@abdn.ac.uk

Dr Robert Bos  
Executive Secretary  
WHO/FAO/UNEP/UNCHS/Panel of  
Experts on Environmental Management  
for Vector Control  
World Health Organisation

CH-1211 Geneva  
SWITZERLAND  
Tel: +41 22 791 35555  
Fax: +41 22 791 4159  
E-mail Bos@WHO.CH

Dr Frank Collins  
Chief, Vector Genetics Section  
Division of Parasitic Diseases  
Centers for Disease Control  
ATLANTA, GEORGIA  
U.S.A.  
Tel: 1-770 488 7465  
Fax: 1-770 488 7794  
e-mail: fhc1@ciddpd2.em.cdc.gov

Prof. Peter Esbjerg  
Chairman, ICIPE Programme Committee  
Royal Vet. and Agricultural University  
Bulowsvej 13, Frederiksberg  
DK-1870 COPENHAGEN  
Denmark  
Tel: 35 35 28 26 86  
Fax: 45 35 28 26 70

Dr Didier Fontenille  
Medical Entomologist  
Head, ORSTOM Laboratory of Medical  
Zoology  
Institut Pasteur  
B.P. 1386, DAKAR  
Senegal  
Tel. 221 32 43 07  
Fax: 221 23 48 74  
e-mail: Fontenil@dakar.orstom.sn

Dr Andrew K. Githeko  
Senior Research Officer (Medical  
Entomologist)  
Kenya Medical Research Institute  
Vector Biology and Control Research  
Centre  
P.O. Box 1578  
Kisumu, KENYA  
Tel. 254-035-44366/7

Dr Richard H. Hunt  
Consultant, Department of Medical  
Entomology  
South African Institute for Medical  
Research  
Hospital Street, P.O. Box 1038  
JOHANNESBURG, 2000  
South Africa  
Tel: 489 93 90  
Fax: 489 90 01

Dr William A. Hawley  
Kenya Medical Research Institute/CDC  
P.O. Box 30137  
NAIROBI,  
Kenya  
Tel: 254-2-713008  
Fax: 254-2-714745  
E-mail: hawley@arcc.or.ke

Dr Flemming Konradsen  
Environment Health Specialist  
International Irrigation Management  
Institute  
P.O. Box 2075  
127, Sunil Mawatha, Battaramulla  
Colombo, SRI LANKA  
Tel: 94 1 86 74 04  
Fax: 94 1 86 68 54  
E-mail: immi@cgnnet.com

Dr Steve Lindsay  
Malaria Unit  
Department for Vector/Biology  
Transmission  
Danish Bilharziasis Laboratory  
Jaegersborg Allé 1D  
DK-2920 Charlottenlund  
DENMARK  
Tel: 5 39 62 61 68  
Fax: 45 39 62 61 21  
E-mail: biladblp@pop.denet.dk

Dr Caroline Maxwell  
National Institute of Medical Research  
Ubwari Field Station  
P.O. Box 81  
Muheza, Mkuo waTanga  
TANZANIA  
Fax: 0873 131 5766

Dr F.W. Moshia  
Director  
Tropical Pesticides Research Institute  
(TPRI)  
P.O. Box 3024  
Arusha  
TANZANIA  
Tel: 057 8813/14  
Fax: 057 8217

Professor Richard Mwangi  
Professor of Zoology  
University of Nairobi  
P.O. Box 30772  
NAIROBI, Kenya  
Tel: 254-2-43185

Dr Graham Reid  
Senior Program Specialist  
Ecosystem Health  
International Development Research  
Centre  
P.O. Box 8500  
OTTAWA, Ontario  
Canada  
Tel: 613 236 6163  
Fax: 1-613 567 7748  
e-mail: greid@idrc.ca

Dr Vincent Robert  
Medical Entomologist  
ORSTOM Laboratory of Medical Zoology  
Institut Pasteur  
DAKAR, Senegal  
Tel: 221 32 09 62  
Fax: 221 23 48 74  
e-mail: robert@belair.orstom.sn

Dr (Mrs.) Sarala K. Subbarao  
Deputy Director  
Malaria Research Centre  
Indian Council of Medical Research  
22 Sham Nath Marg  
DELHI 110054  
INDIA  
Tel: 91 11 235 721/224 7983  
Fax: 91 11 72 34 234  
e-mail: vps@mrcicmr.ren.nic.in

Dr Willem Takken  
Medical/Veterinary Entomologist  
Wageningen Agricultural University  
Department of Entomology  
THE NETHERLANDS  
Tel: 31 8370 484 652  
Fax: 31 8370 84821  
e-mail:willem  
takken@medew.ento.wau.nl

Dr Fantamady Sekou Traore  
Medical Entomologist (Cytognitist)  
Malaria and Training Centre  
National School of Medicine and  
Pharmacy  
Research Pharmacie BP 1805  
MALI  
Tel: 223 22 52 77  
Fax: 223 22 98 79  
e-mail: cheick@mrtcbko.malinet.ml

#### ICIPE STAFF

Dr. Hans R. Herren, *Director General*  
Prof. Ahmed Hassanali, *Interim Deputy Director General*  
Mrs. Rhoda A. Odingo, *Director International Cooperation*  
Dr. Steve Mihok, *Ag. Programme Leader, Disease Vector Management Programme*  
Dr. E.O. Osir, *Head, Molecular Biology and Biochemistry Department*  
Dr. N.K. Maniania, *Ag. Head, Pathology and Microbiology Department*  
Prof. Fassil G. Kiros, *Head, Social Science Department*  
Dr. K. Ampong-Nyarko, *Ag. Head, Ecology and Biodiversity Department*  
Dr. R.K. Saini, *Ag. Head, Behavioural Biology Department*  
Dr. L. Lwande, *Ag. Head, Chemical Ecology Department*  
Dr. C.M. Mutero, *EU Project Coordinator*  
Dr. Muhinda E. Mugunga, *ICIPE/Hebrew Collaborative Project*



**ICIPE Science Press**  
PO Box 72913  
Nairobi, Kenya